

For Research Use Only

IKZF1 Polyclonal antibody

Catalog Number: 12016-1-AP **7 Publications**



Basic Information

Catalog Number: 12016-1-AP	GenBank Accession Number: BC018349	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 µg/ml by Nanodrop;	GeneID (NCBI): 10320	Recommended Dilutions: WB 1:500-1:3000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500
Source: Rabbit	Full Name: IKAROS family zinc finger 1 (Ikaros)	
Isotype: IgG	Calculated MW: 477 aa, 53 kDa	
Immunogen Catalog Number: AG2631	Observed MW: 50-70 kDa	

Applications

Tested Applications: IHC, IP, WB, ELISA	Positive Controls: WB : HeLa cells, HepG2 cells IP : HeLa cells, IHC : human lymphoma tissue,
Cited Applications: CoIP, IHC, WB	
Species Specificity: human	
Cited Species: human, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

IKZF1, also known as DNA-binding protein Ikaros, belongs to the Ikaros C2H2-type zinc-finger protein family. The N-terminal zinc-fingers 2 and 3 are required for DNA binding as well as for targeting IKZF1 to pericentromeric heterochromatin. The C-terminal zinc-finger domain is required for dimerization. IKZF1 is abundantly expressed in thymus, spleen and peripheral blood leukocytes and lymph nodes. IKZF1 is localized in nucleus. IKZF1 as a transcription regulator of hematopoietic cell differentiation binds gamma-satellite DNA and plays a role in the development of lymphocytes, B- and T-cells. Binds and activates the enhancer (delta-A element) of the CD3-delta gene. IKZF1 is a repressor of the TDT (fikzfterminal deoxynucleotidyltransferase) gene during thymocyte differentiation. IKZF1 exists some isoforms with 50-70 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Konstantinos Dimopoulos	29718735	Leuk Lymphoma	IHC
Morgann Klink	33807974	Cancers (Basel)	WB
Yan Sun	28698179	Circ Res	CoIP

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using 12016-1-AP (IKZF1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-IKZF1 (IP:12016-1-AP, 4ug; Detection:12016-1-AP 1:500) with HeLa cells lysate 1200ug.



HeLa cells were subjected to SDS PAGE followed by western blot with 12016-1-AP (IKZF1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.