

For Research Use Only

SLU7 Polyclonal antibody

Catalog Number: 12050-1-AP **2 Publications**



Basic Information

Catalog Number: 12050-1-AP	GenBank Accession Number: BC010634	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 300 µg/ml by Nanodrop and 133 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 10569	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: O95391	
Isotype: IgG	Full Name: SLU7 splicing factor homolog (S. cerevisiae)	
Immunogen Catalog Number: AG2680	Calculated MW: 586 aa, 68 kDa	
	Observed MW: 68 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : Jurkat cells, HeLa cells, mouse colon tissue, K-562 cells, Y79 cells
Cited Applications: WB, IF, RIP, IHC	IHC : mouse pancreas tissue, human colon cancer tissue, rat pancreas tissue, rat stomach tissue
Species Specificity: human, mouse, rat	
Cited Species: human, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Pre-mRNA splicing occurs in two sequential transesterification steps. SLU7 is a splicing factor that has been found to be essential during the second catalytic step in the pre-mRNA splicing process. It associates with the spliceosome and contains a zinc knuckle motif that is found in other splicing factors and is involved in protein-nucleic acid and protein-protein interactions.

Notable Publications

Author	Pubmed ID	Journal	Application
Xianchong Zheng	38084438	Adv Sci (Weinh)	WB,IHC,IF
Runjie Song	37408008	Mol Cancer	RIP

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 12050-1-AP (SLU7 antibody) at dilution of 1:100 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human colon cancer using 12050-1-AP (SLU7 antibody) at dilution of 1:50 (under 10x lens).




Various lysates were subjected to SDS PAGE followed by western blot with 12050-1-AP (SLU7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 12050-1-AP (SLU7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 12050-1-AP (SLU7 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat pancreas tissue slide using 12050-1-AP (SLU7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat stomach tissue slide using 12050-1-AP (SLU7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).