For Research Use Only

PRKCSH Polyclonal antibody

Catalog Number: 12148-1-AP

Featured Product

8 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12148-1-AP BC013586

GeneID (NCBI): Size

150ul, Concentration: 1200 µg/ml by 5589 Nanodrop and 533 $\mu g/ml$ by Bradford Full Name:

method using BSA as the standard; protein kinase C substrate 80K-H

Calculated MW: Rabbit 80 kDa Isotype: Observed MW:

IgG 80 kDa

Immunogen Catalog Number:

AG2796

human, rat, mouse

Purification Method: Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF 1:50-1:500

WB 1:2000-1:8000

Applications

Tested Applications: Positive Controls:

IF, IHC, IP, WB, ELISA WB: A431 cells, mouse liver tissue, rat liver tissue, **Cited Applications:** HeLa cells, Jurkat cells

IF, IHC, WB IP: HeLa cells,

Species Specificity: IHC: human kidney tissue, human, mouse, rat

IF: HeLa cells, **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

 $PRKCSH\ encodes\ the\ beta-subunit\ of\ glucosidase\ II,\ an\ N-linked\ glycan-processing\ enzyme\ in\ the\ endoplasmic$ reticulum (ER). This protein is an acidic phospho-protein known to be a substrate for protein kinase C. Defects in PRKCSH are a cause of an autosomal dominant polycystic liver disease (PCLD). Glucosidase II is an ER-localized enzyme that contains a and β subunits (Glucosidase IIa and Glucosidase IIB) which form a defined heterodimeric complex.

Notable Publications

Author	Pubmed ID	Journal	Application
Ling Leng	35679865	Cell Rep	IHC
Birong Liang	35598104	Exp Physiol	IHC
Gu-Choul Shin	31320625	Nat Commun	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12148-1-AP (PRKCSH antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




IP Result of anti-PRKCSH (IP:12148-1-AP, 4ug; Detection:12148-1-AP 1:1000) with HeLa cells lysate 2000ug.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 12148-1-AP (PRKCSH antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 12148-1-AP (PRKCSH antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 12148-1-AP (PRKCSH antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).