For Research Use Only

TRIP4 Polyclonal antibody

Catalog Number: 12324-1-AP 1 Publications



Purification Method:

WB 1:500-1:2000

IHC 1:50-1:500

WB: HL-60 cells, COLO 320 cells, HeLa cells

IHC: human liver cancer tissue,

Positive Controls:

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

12324-1-AP BC012448
Size: GeneID (NCBI):
150ul , Concentration: 500 ug/ml by 9325

Nanodrop; UNIPROT ID:
Source: Q15650
Rabbit Full Name:

Isotype: thyroid hormone receptor interactor 4

IgG Calculated MW:
Immunogen Catalog Number: 581 aa, 66 kDa
AG2987 Observed MW:
60-65 kDa

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Thyroid hormone receptor interactor 4(TRIP4), other named ASC1, is a transcriptional coactivator that promote transcriptional efficiencies. It harbors an autonomous transactivation domain that contains a putative zinc finger motif, which serves as a binding site for nuclear receptors for thyroid, estrogen, all-trans-retinoic acid and RXR by interaction with SRC-1 or CBP/p300. There are two forms of ASC1 have been detected in mRNA level, isoform1(581aa, ~65kd) and isoform2(539aa, ~60kd)(12077347,12390891)

Notable Publications

Author	Pubmed ID	Journal	Application
Min Ni	33473208	Genet Med	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HL-60 cells were subjected to SDS PAGE followed by western blot with 12324-1-AP (TRIP4 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 12324-1-AP (TRIP4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 12324-1-AP (TRIP4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 12324-1-AP (TRIP4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 12324-1-AP (TRIP4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).