For Research Use Only

EBI3 Polyclonal antibody

Catalog Number: 12371-1-AP

Featured Product

3 Publications

BC015364

25-35 kDa



Basic Information

Catalog Number: GenBank Accession Number: 12371-1-AP

GeneID (NCBI):

150ul, Concentration: 300 µg/ml by 10148

Nanodrop and 233 µg/ml by Bradford Full Name: method using BSA as the standard; Epstein-Barr virus induced 3

Calculated MW: Rabbit 229 aa, 25 kDa Isotype: Observed MW:

Immunogen Catalog Number:

AG3051

IgG

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB IHC 1:50-1:500

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

IF. WB

Species Specificity: human, pig **Cited Species:**

human, mouse Note-IHC: suggested antigen retrieval with

retrieval may be performed with citrate

TE buffer pH 9.0; (*) Alternatively, antigen

buffer pH 6.0

Positive Controls:

WB: human placenta tissue, pig spleen tissue

IP: human placenta tissue,

IHC: human spleen tissue, human placenta tissue

Background Information

EBI3, a 34-kDa secreted glycoprotein, is a subunit in 2 distinct heterodimeric cytokines: IL-27 and IL-35. IL-27, $composed \ of \ p28 \ (IL-27) \ and \ EBI3, can \ trigger \ signaling \ in \ T \ cells, \ B \ cells, \ and \ myeloid \ cells. \ IL-35, \ composed \ of \ EBI3$ and the ubiquitously expressed p35 subunit of IL-12, is an inhibitory cytokine involved in regulatory T-cell function. EBI3 plays a role in regulating cell-mediated immune responses. (PMID: 8551575; 18033300; 17874423; 21931777)

Notable Publications

Author	Pubmed ID	Journal	Application
Qi Zuo	34145061	J Immunol	WB
Tong Zhang	36548354	PLoS One	WB,IF
Jia-Hui Deng	30753412	Pathog Dis	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




human placenta tissue was subjected to SDS PAGE followed by western blot with 12371-1-AP (EBI3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-EBI3(IP:12371-1-AP, 4ug; Detection:12371-1-AP 1:800) with human placenta tissue lysate 600 ug.



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 12371-1-AP (EBI3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 12371-1-AP (EBI3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).