For Research Use Only

EIF3C Polyclonal antibody

Catalog Number: 12733-1-AP 1 Publications



Purification Method:

WB 1:500-1:1000

protein lysate

IHC 1:50-1:500

IHC: human stomach cancer tissue, human colon

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

12733-1-AP BC001571 GeneID (NCBI): Size:

150ul, Concentration: 650 ug/ml by 8663 Nanodrop and 400 ug/ml by Bradford UNIPROT ID: method using BSA as the standard;

Q99613 Source: Full Name:

Rabbit eukaryotic translation initiation factor 3, subunit C Isotype:

IgG Calculated MW: 913 aa, 110 kDa Immunogen Catalog Number: AG3438 Observed MW:

110 kDa

Applications

Tested Applications: WB, IHC, IP, ELISA

Cited Applications:

Species Specificity: human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Author Yanan Pi **Pubmed ID** 37828084

Journal Cell Death Differ

Positive Controls:

IP: K-562 cells,

cancer tissue

WB: K-562 cells, MCF-7 cells

Application WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

Notable Publications


Selected Validation Data



K-562 cells were subjected to SDS PAGE followed by western blot with 12733-1-AP (EIF3C antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-EIF3C (IP:12733-1-AP, 4ug; Detection:12733-1-AP 1:500) with K-562 cells lysate 3200ug.



Immunohistochemical analysis of paraffinembedded stomach cancer slide using 12733-1-AP (EIF3C antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).