

For Research Use Only

SEC5/EXOC2 Polyclonal antibody

Catalog Number: 12751-1-AP

Featured Product

20 Publications



Basic Information

Catalog Number:

12751-1-AP

Size:

150ul, Concentration: 260 ug/ml by Nanodrop and 233 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3147

GenBank Accession Number:

BC016918

GeneID (NCBI):

55770

UNIPROT ID:

Q96KP1

Full Name:

exocyst complex component 2

Calculated MW:

924 aa, 104 kDa

Observed MW:

95-100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, human brain tissue, human ileum tissue

IP: mouse brain tissue,

IHC: human breast cancer tissue,

IF/ICC: HeLa cells,

Background Information

EXOC2 (exocyst complex component 2), also known as SEC5 and SEC5L1, is a component of the exocyst complex, and is required to mediate RalB-dependent survival signals in transformed cells. The exocyst complex, composed of eight evolutionarily conserved subunits (SEC3, SEC5, SEC6, SEC8, SEC10, SEC15, EXO70, and EXO84), is involved in tethering post-Golgi secretory vesicles to specific plasma membrane domains. The gene of EXOC2 maps to chromosome 6p25.3, and encodes a 924-amino acid protein with an experimentally determined molecular mass of 95-100 kDa. EXOC2 mRNA is widely expressed with highest levels in brain and placenta.

Notable Publications

Author	Pubmed ID	Journal	Application
Sánchez-Ruiz Jesús J	21810610	J Immunol	WB
Hiroki Takeuchi	26617273	Cell Microbiol	WB
Maeran Uhm	28325821	Sci Signal	WB, CoIP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 12751-1-AP (SEC5/EXOC2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-SEC5/EXOC2 (IP:12751-1-AP, 5ug; Detection:12751-1-AP 1:500) with mouse brain tissue lysate 3000ug.



Immunohistochemical analysis of paraffin-embedded human breast cancer using 12751-1-AP (SEC5/EXOC2 antibody) at dilution of 1:100 (under 10x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 12751-1-AP (SEC5/EXOC2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).