## For Research Use Only

## MLLT3/AF9 Polyclonal antibody

Catalog Number: 12825-1-AP



**Purification Method:** 

WB 1:500-1:1000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number:

12825-1-AP BC036089 GeneID (NCBI):

150ul , Concentration: 500 ug/ml by Nanodrop and 400 ug/ml by Bradford  $\,$  UNIPROT ID:

method using BSA as the standard; Source:

Full Name: Rabbit myeloid/lymphoid or mixed-lineage

leukemia (trithorax homolog, Isotype Drosophila); translocated to, 3

Calculated MW: Immunogen Catalog Number: AG3528 568 aa, 63 kDa Observed MW:

63-70 kDa

P42568

**Applications** 

**Tested Applications:** 

WB. FIISA

Species Specificity: human, mouse, rat

**Positive Controls:** 

WB: Jurkat cells, HL-60 cells

## **Background Information**

MLLT3, also named as AF9, YEATS3, is a 568 amino acid protein, which is a component of the superelongation complex6 and co-operates with DOT1L, which di/trimethylates H3K79 to promote transcription. MLLT3 localizes to active transcription start sites (TSSs) through the YEATS domain, which recognizes active histone marks such as H3K9 acetylation and crotonylation. A truncated MLLT3 that lacks the YEATS domain forms a leukaemic fusion protein with the N terminus of MLL1, which misdirects MLLT3-interacting complexes to induce aberrant gene transcription. MLLT3 also regulates erythroid or megakaryocytic progenitors and was identified as a definitive HSC hub gene during mouse development. Researcher found that the expression of a gene called MLLT3 was closely correlated with blood stem cells'potential to self-renew and that the protein generated by the MLLT3 gene provides blood stem cells with the instructions necessary to maintain its ability to self-renew. It does this by working with other regulatory proteins to keep important parts of the blood stem cell's machinery operational as the cells divide.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 12825-1-AP (MLLT3/AF9 antibody at dilution of 1:600 incubated at room temperature for 1.5 hours.