For Research Use Only

DTL Polyclonal antibody

Catalog Number: 12896-1-AP

2 Publications



Basic Information

Catalog Number: 12896-1-AP

GenBank Accession Number: BC033540

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 200 µg/ml by

51514

WB 1:200-1:1000 IP 0.5-4.0 ug for IP and 1:500-1:1000

Nanodrop and 133 µg/ml by Bradford Full Name:

denticleless homolog (Drosophila)

for WB

method using BSA as the standard;

Calculated MW:

730 aa, 79 kDa Observed MW:

79 kDa

IgG

Rabbit

Isotype:

Immunogen Catalog Number:

AG3589

IF 1:50-1:500

Applications

Tested Applications: IF, IP, WB, ELISA

WB: HeLa cells, LO2 cells, MCF-7 cells

Cited Applications:

IP: mouse testis tissue,

Positive Controls:

IF: HeLa cells,

IF, WB

Species Specificity: human, mouse, rat

Cited Species:

human, mouse

Background Information

CDT2, also named as L2DTL or RAMP, is one of the substrate receptors of the Cullin Ring Ubiquitin Ligase 4 that targets for ubiquitin mediated degradation a number of substrates, such as CDT1, p21 and CHK1, involved in the regulation of cell cycle and survival. CDT2 overexpression was reported in breast (PMID: 18542055), gastric (PMID: 19672268) and ovarian carcinomas (PMID: 23995842) and rhabdomy osarcomas (PMID: 19235922) and associated with the aggressiveness of hepatocellular carcinomas (PMID: 17106265).

Notable Publications

Author	Pubmed ID	Journal	Application
Chunxia Zhang	35108058	Sci Adv	IF
Chengli Yu	36843329	Proteomics	WB

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 12896-1-AP (DTL antibody) at dilution of 1:200 incubated at room temperature for 1.5 hours.



IP Result of anti-DTL (IP:12896-1-AP, 3ug; Detection:12896-1-AP 1:500) with mouse testis tissue lysate 8000ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 12896-1-AP (DTL antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).