### For Research Use Only

# TRIM69 Polyclonal antibody

Catalog Number: 12951-1-AP

**Featured Product** 

6 Publications



**Purification Method:** 

WB 1:500-1:1000 IHC 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

**Applications** 

Catalog Number: GenBank Accession Number:

12951-1-AP BC033314

Size: GeneID (NCBI): 150ul , Concentration: 800 µg/ml by 140691

Nanodrop; Full Name:
Source: tripartite motif-containing 69

Rabbit Calculated MW:
Isotype: 500 aa, 57 kDa
IgG Observed MW:
Immunogen Catalog Number: 60-70 kDa

AG4074

Tested Applications: Po

IHC, WB, ELISA
Cited Applications:
CoIP, IF, IHC, IP, WB
Species Specificity:
human, mouse, rat
Cited Species:

human, sheep, mouse, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

. .

Positive Controls:

WB: mouse testis tissue, HeLa cells IHC: human gliomas tissue,

# **Background Information**

E3 ubiquitin-protein ligase TRIM69 (TRIM69) plays an important role in antiviral immunity by restricting different viral infections including dengue virus or vesicular stomatitis Indiana virus (PubMed:31375575, PubMed:31578292). It plays also a role in cataract formation together with TP53 (PubMed:30844644). Mechanistically, inhibits UVB-induced cell apoptosis and reactive oxygen species (ROS) production by inducing TP53 ubiquitination (PubMed:30844644). It has 4 isoforms with molecular weights of 57, 39, 34, and 32 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Huiyuan Jing	29123520	Front Immunol	WB
Tonya Kueck	31578292	J Virol	WB
Zhongjia Jiang	28553017	Mediators Inflamm	WB

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## **Selected Validation Data**



mouse testis tissue were subjected to SDS PAGE followed by western blot with 12951-1-AP (TRIM69 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas using 12951-1-AP (TRIM69 antibody) at dilution of 1:100 (under 10x lens).