For Research Use Only

PPM1A Polyclonal antibody Catalog Number:12961-1-AP 3 Publications




Basic Information	Catalog Number: 12961-1-AP	ord UNIPROT ID:		Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:8000 IHC 1:500-1:2000					
	Size:								
	150ul , Concentration: 900 ug/ml by								
	Nanodrop and 333 ug/ml by Bradford method using BSA as the standard;								
	Source:	Full Name:							
	Rabbit	protein phosphatase 1A (formerly 2C),							
	Isotype: IgG	magnesium-dependent, alpha isoform							
	Immunogen Catalog Number: AG3645	Calculated MW: 382 aa, 42 kDa							
		Observed MW 42 kDa							
Applications	Tested Applications:	Positive Controls:							
	WB, IHC, ELISA			WB : Jurkat cells, HepG2 cells, human liver tissue, rat kidney tissue, mouse kidney tissue, K-562 cells IHC : human breast cancer tissue, human urothelial carcinoma tissue					
	Cited Applications: WB, IF		-						
	Species Specificity:								
	human, mouse, rat Cited Species: mouse, rat Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0								
					Background Information	PPM1A(Protein phosphatase 1A) is also named as PPPM1A and belongs to the PP2C family. It is a serine/threonine protein phosphatase that is essential for regulating cellular stress responses in eukaryotes. PPM1A plays an important role in tumor suppression and that it negatively controls many signaling pathways which are critical for trophoblast invasion.(PMID:19404668). This protein distributes in both the cytoplasm and nucleus of mammalian cells(PMID:9003755). It has 2 isoforms produced by alternative splicing.			
					Notable Publications	Author Pu	bmed ID	Journal	Application
Haibin Tian 36	185374	iScience	IF						
	491170	Biol Pharm Bull	WB						
Katyayanee Neopane 36	543129	Cell Rep							
itorage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50		7.3.						
	Aliquoting is unnecessary for -20°C s	torage							
** 20ul sizes contain 0.1% BSA									
*** 20ul sizes contain 0.1% BSA									


in USA), or 1(312) 455-8498 (outside USA)

Proteintech@ptg
W: ptglab.com

other manufacturer.


Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 12961-1-AP (PPM1A antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.

Various lysates were subjected to SDS PAGE followed by western blot with 12961-1-AP (PPM1A antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 12961-1-AP (PPM1A antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).