## For Research Use Only

## NKX3-1 Polyclonal antibody

Catalog Number: 13069-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number: 13069-1-AP

BC074864

NK3 homeobox 1

Size: GeneID (NCBI):

150ul, Concentration: 350 µg/ml by 4824

Nanodrop and 200 µg/ml by Bradford Full Name:

method using BSA as the standard;

Calculated MW: Rabbit 234 aa, 26 kDa Isotype: Observed MW: IgG 34-38 kDa

Immunogen Catalog Number:

AG4070

**Purification Method:** Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IHC 1:400-1:1600

**Applications** 

**Tested Applications:** 

IHC, WB, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Positive Controls:** 

WB: LNCaP cells, human testis tissue IHC: human prostate cancer tissue,

## **Background Information**

NKX3-1, also named as NKX3.1 and NKX3A, belongs to the NK-3 homeobox family. It is a transcription factor, which binds preferentially the consensus sequence 5'-TAAGT[AG]-3' and can behave as a transcriptional repressor. NKX3-1 plays an important role in normal prostate development, regulating proliferation of glandular epithelium and in the formation of ducts in prostate. It act as a tumor suppressor controlling prostate carcinogenesis, as shown by the ability to inhibit proliferation and invasion activities of PC-3 prostate cancer cells(PMID:19462257). The calcualted molecular weight of NKX3-1 is 26 kDa, but modified NKX3-1 is about 32-35 kDa. (PMID: 18597829)

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




LNCaP cells were subjected to SDS PAGE followed by western blot with 13069-1-AP (NK/3-1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



human testis tissue were subjected to SDS PAGE followed by western blot with 13069-1-AP (NKX3-1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 13069-1-AP (NKX3-1 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 13069-1-AP (NKX3-1 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).