
For Research Use Only

PACSIN1 Polyclonal antibody Catalog Number:13219-1-AP 2 Publications

proteintech® Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 13219-1-AP	GenBank Accession Number: BC040228	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 850 ug/ml by Nanodrop and 393 ug/ml by Bradford method using BSA as the standard;	29993	WB 1:500-1:2000	
		UNIPROT ID: Q9BY11	IHC 1:50-1:500 IF/ICC 1:20-1:200	
	Source: Rabbit	Full Name: protein kinase C and casein kinase substrate in neurons 1 Calculated MW: 444 aa, 51 kDa		
	Isotype: IgG			
	Immunogen Catalog Number: AG4102			
		Observed MW: 48-51 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, ELISA	WB : mouse	WB: mouse cerebellum tissue, mouse brain tissue	
	Cited Applications: WB, IHC	IHC : human	brain tissue,	
	Species Specificity: human, mouse, rat	IF/ICC : SH-SY5Y cells,		
	Cited Species: human, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	(PACSIN) family. In mammals, the PA (PMID: 34990060). PACSIN1 is express tissues, and PACSIN3 is expressed ma members contain an N-terminal F-BA and endosomal recycling. Meanwhile particular role in membrane shaping neuromorphogenesis and the regulat	ACSIN family is comprised of three m sed mainly in neurons, whereas PAC ainly in skeletal muscle and the hea R domain and a C-terminal SH3 dom e, it has a role in actin remodeling an and reconstruction (PMID: 23035120 ion of the nervous system, and the ir	SIN2 is ubiquitously expressed in all rt (PMID: 23668323). All of these three ain. PACSIN1 plays a role in endocytosi: d microtubule nucleation and also plays 34422904). PACSIN1 is involved in appropriate expression of PACSIN1 has	
	(PACSIN) family. In mammals, the PA (PMID: 34990060). PACSIN1 is express tissues, and PACSIN3 is expressed ma members contain an N-terminal F-BA and endosomal recycling. Meanwhile particular role in membrane shaping neuromorphogenesis and the regulat been associated with some neurologi disease (PMID: 34990060).	ACSIN family is comprised of three m sed mainly in neurons, whereas PAC ainly in skeletal muscle and the hea R domain and a C-terminal SH3 dom e, it has a role in actin remodeling an and reconstruction (PMID: 23035120 ion of the nervous system, and the in ical diseases, including schizophreni	embers, PACSIN1, PACSIN2, and PACSIN SIN2 is ubiquitously expressed in all rt (PMID: 23668323). All of these three ain. PACSIN1 plays a role in endocytosis d microtubule nucleation and also plays 34422904). PACSIN1 is involved in appropriate expression of PACSIN1 has a, Alzheimer's disease, and Huntington's	
	(PACSIN) family. In mammals, the PA (PMID: 34990060). PACSIN1 is express tissues, and PACSIN3 is expressed ma members contain an N-terminal F-BA and endosomal recycling. Meanwhile particular role in membrane shaping neuromorphogenesis and the regulat been associated with some neurologi disease (PMID: 34990060). Author Put	ACSIN family is comprised of three m sed mainly in neurons, whereas PAC ainly in skeletal muscle and the hea R domain and a C-terminal SH3 dom e, it has a role in actin remodeling an and reconstruction (PMID: 23035120) ion of the nervous system, and the in ical diseases, including schizophreni	embers, PACSIN1, PACSIN2, and PACSIN SIN2 is ubiquitously expressed in all rt (PMID: 23668323). All of these three ain. PACSIN1 plays a role in endocytosis d microtubule nucleation and also plays 34422904). PACSIN1 is involved in appropriate expression of PACSIN1 has a, Alzheimer's disease, and Huntington's Application	
	(PACSIN) family. In mammals, the PA (PMID: 34990060). PACSIN1 is express tissues, and PACSIN3 is expressed mi members contain an N-terminal F-BA and endosomal recycling. Meanwhile particular role in membrane shaping neuromorphogenesis and the regulat been associated with some neurologi disease (PMID: 34990060). Author Put Zhou Zimu 344	ACSIN family is comprised of three m sed mainly in neurons, whereas PAC ainly in skeletal muscle and the hea R domain and a C-terminal SH3 dom e, it has a role in actin remodeling an and reconstruction (PMID: 23035120 ion of the nervous system, and the in ical diseases, including schizophreni cal diseases, Front Mol Biosci	Ambers, PACSIN1, PACSIN2, and PACSIN SIN2 is ubiquitously expressed in all rt (PMID: 23668323). All of these three ain. PACSIN1 plays a role in endocytosis d microtubule nucleation and also plays 34422904). PACSIN1 is involved in appropriate expression of PACSIN1 has a, Alzheimer's disease, and Huntington's Application IHC	
	(PACSIN) family. In mammals, the PA (PMID: 34990060). PACSIN1 is express tissues, and PACSIN3 is expressed mi members contain an N-terminal F-BA and endosomal recycling. Meanwhile particular role in membrane shaping neuromorphogenesis and the regulat been associated with some neurologi disease (PMID: 34990060). Author Put Zhou Zimu 344	ACSIN family is comprised of three m sed mainly in neurons, whereas PAC ainly in skeletal muscle and the hea R domain and a C-terminal SH3 dom e, it has a role in actin remodeling an and reconstruction (PMID: 23035120) ion of the nervous system, and the in ical diseases, including schizophreni	embers, PACSIN1, PACSIN2, and PACSIN SIN2 is ubiquitously expressed in all rt (PMID: 23668323). All of these three ain. PACSIN1 plays a role in endocytosid d microtubule nucleation and also plays 34422904). PACSIN1 is involved in appropriate expression of PACSIN1 has a, Alzheimer's disease, and Huntington's Application	
Background Information	(PACSIN) family. In mammals, the PA (PMID: 34990060). PACSIN1 is express tissues, and PACSIN3 is expressed mammbers contain an N-terminal F-BA and endosomal recycling. Meanwhile particular role in membrane shaping neuromorphogenesis and the regulat been associated with some neurologi disease (PMID: 34990060). Author Put Zhou Zimu 344 Shaoran Shen 393 Storage: Storage star - 20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50	ACSIN family is comprised of three m sed mainly in neurons, whereas PAC ainly in skeletal muscle and the hea R domain and a C-terminal SH3 dom e, it has a role in actin remodeling an and reconstruction (PMID: 23035120 ion of the nervous system, and the in ical diseases, including schizophreni actual diseases, including schizophreni actual seases, including schizophreni ac	embers, PACSIN1, PACSIN2, and PACSIN SIN2 is ubiquitously expressed in all rt (PMID: 23668323). All of these three ain. PACSIN1 plays a role in endocytosis d microtubule nucleation and also plays 34422904). PACSIN1 is involved in appropriate expression of PACSIN1 has a, Alzheimer's disease, and Huntington's Application IHC	
Notable Publications	(PACSIN) family. In mammals, the PA (PMID: 34990060). PACSIN1 is express tissues, and PACSIN3 is expressed mammbers contain an N-terminal F-BA and endosomal recycling. Meanwhile particular role in membrane shaping neuromorphogenesis and the regulat been associated with some neurologi disease (PMID: 34990060). Author Put Zhou Zimu 344 Shaoran Shen 393 Storage: Storage: Storage Buffer:	ACSIN family is comprised of three m sed mainly in neurons, whereas PAC ainly in skeletal muscle and the hea R domain and a C-terminal SH3 dom e, it has a role in actin remodeling an and reconstruction (PMID: 23035120 ion of the nervous system, and the in ical diseases, including schizophreni actual diseases, including schizophreni actual seases, including schizophreni ac	embers, PACSIN1, PACSIN2, and PACSIN SIN2 is ubiquitously expressed in all rt (PMID: 23668323). All of these three ain. PACSIN1 plays a role in endocytosi. d microtubule nucleation and also plays 34422904). PACSIN1 is involved in appropriate expression of PACSIN1 has a, Alzheimer's disease, and Huntington's Application IHC	


Selected Validation Data




mouse cerebellum tissue were subjected to SDS PAGE followed by western blot with 13219-1-AP (PACSIN1 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 13219-1-AP (PACSIN1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 13219-1-AP (PACSIN1 Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of SH-SY5Y cells using 13219-1-AP (PACSIN1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).