#### For Research Use Only

# PHKB Polyclonal antibody

Catalog Number: 13400-1-AP

**Featured Product** 

4 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

13400-1-AP BC033657
Size: Genel D (NCBI):

150ul, Concentration: 450 ug/ml by 5257

Nanodrop; UNIPROT ID:
Source: Q93100
Rabbit Full Name:

lsotype: phosphorylase kinase, beta

IgG Calculated MW:
Immunogen Catalog Number: 1086 aa, 124 kDa
AG4240 Observed MW:

124 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

## Applications

Tested Applications:
WB, IP, IHC, ELISA
Cited Applications:
WB, IHC, IP
Species Specificity:
human, mouse
Cited Species:

human, mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: HepG2 cells, Jurkat cells, K-562 cells, LNCaP cells, mouse heart tissue, rat heart tissue, rat skeletal muscle tissue, mouse skeletal muscle tissue

IP: mouse heart tissue.

IHC: mouse liver tissue, human skeletal muscle tissue, human liver tissue, mouse skeletal muscle tissue

# **Background Information**

PHKB gene encodes phosphorylase kinase subunit beta involved in glycan biosynthesis and glycogen metabolism. PHKB activity is regulated by phosphorylation of various serine residues, and catalyzes the phosphorylation of serine in certain substrates, including troponin I. Phosphorylase kinase (PhK) complex, composed of alpha, beta, gamma, and delta subunits, stimulates energy production from glycogen in the cascade activation of glycogenolysis. Its large homologous alpha and beta subunits regulate the activity of the catalytic gamma subunit. Defects in PHKB are the cause of glycogen storage disease type 9B (GSD9B) also known as phosphorylase kinase deficiency of liver and muscle (PKD), characterized by hepathomegaly, only slightly elevated transaminases and plasma lipids, clinical improvement with increasing age, and remarkably no clinical muscle involvement.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Guanghui Wang	28275865	J Cancer Res Clin Oncol	WB,IHC
Motoyasu Hosokawa	30870781	iScience	WB
Haigang Cao	38823637	J Biol Chem	WB

### Storage

Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


# **Selected Validation Data**




HepG2 cells were subjected to SDS PAGE followed by western blot with 13400-1-AP (PHKB antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 13400-1-AP (PHKB antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP result of anti-PHKB (IP:13400-1-AP, 4ug; Detection:13400-1-AP 1:500) with mouse heart tissue lysate 1680 ug.




Immunohistochemical analysis of paraffinembedded human skeletal muscle tissue slide using 13400-1-AP (PHKB Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 13400-1-AP (PHKB antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 13400-1-AP (PHKB antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human skeletal muscle tissue slide using 13400-1-AP (PHKB Antibody) at dilution of 1:50 (under 40x lens).