For Research Use Only

AUP1 Polyclonal antibody Catalog Number: 13726-1-AP 6 Publications

oroteintech Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 13726-1-AP	GenBank Accession Number: BC033646	Purification Method: Antigen affinity purification	
	Size: 150ul, Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG4680	GeneID (NCBI): 550 UNIPROT ID: Q9Y679 Full Name: ancient ubiquitous protein 1 Calculated MW: 416 aa, 45 kDa Observed MW: 40 kDa	Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200 IF/ICC 1:10-1:100	
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, IP, ELISA Cited Applications:	WB : K-56	52 cells, HEK-293 cells, Jurkat cells	
	WB, IF	IP : HEK-293 cells,		
	Species Specificity:		IHC : human liver cancer tissue, IF/ICC : A431 cells,	
	human, mouse, rat Cited Species:	IF/ICC:/	4431 Cells,	
	human			
	Note-IHC: suggested antigen (TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen		
Background Information	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 AUP1 (Ancient ubiquitous protein 1) surface of ER and Lipid droplets, invo 37029364). In addition, AUP1 also co	vely, antigen with citrate is a protein associated with lipid r lved in the degradation of misfold ntrols lipid synthesis as it induces s of lipid biosynthesis, such as the	ded proteins through autophagy (PMID: subiquitination and the subsequent	
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 AUP1 (Ancient ubiquitous protein 1) surface of ER and Lipid droplets, invo 37029364). In addition, AUP1 also co degradation of several key regulator 3-methylglutaryl-coenzyme A reduc	vely, antigen with citrate is a protein associated with lipid r lved in the degradation of misfold ntrols lipid synthesis as it induces s of lipid biosynthesis, such as the	ded proteins through autophagy (PMID: subiquitination and the subsequent	
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 AUP1 (Ancient ubiquitous protein 1) surface of ER and Lipid droplets, invo 37029364). In addition, AUP1 also co degradation of several key regulator 3-methylglutaryl-coenzyme A reduct Author Put	vely, antigen with citrate is a protein associated with lipid r lved in the degradation of misfold ntrols lipid synthesis as it induces s of lipid biosynthesis, such as the ase (PMID: 28330944).	ded proteins through autophagy (PMID: subiquitination and the subsequent e cholesterol biosynthetic enzyme 3-hydroxy	
Background Information Notable Publications	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 AUP1 (Ancient ubiquitous protein 1) surface of ER and Lipid droplets, invo 37029364). In addition, AUP1 also co degradation of several key regulator 3-methylglutaryl-coenzyme A reduct Author Pu Milton To 27	vely, antigen with citrate is a protein associated with lipid r lved in the degradation of misfold ntrols lipid synthesis as it induces s of lipid biosynthesis, such as the case (PMID: 28330944).	s ubiquitination and the subsequent e cholesterol biosynthetic enzyme 3-hydroxy Application	
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 AUP1 (Ancient ubiquitous protein 1) surface of ER and Lipid droplets, invo 37029364). In addition, AUP1 also co degradation of several key regulator 3-methylglutaryl-coenzyme A reduct Author Pu Milton To 27 Edmond Y Huang 29	vely, antigen with citrate	ded proteins through autophagy (PMID: subiquitination and the subsequent e cholesterol biosynthetic enzyme 3-hydroxy Application WB	
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 AUP1 (Ancient ubiquitous protein 1) surface of ER and Lipid droplets, invo 37029364). In addition, AUP1 also co degradation of several key regulator 3-methylglutaryl-coenzyme A reduct Author Pu Milton To 27 Edmond Y Huang 29	vely, antigen vith citrate	ded proteins through autophagy (PMID: subiquitination and the subsequent e cholesterol biosynthetic enzyme 3-hydroxy Application WB	
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 AUP1 (Ancient ubiquitous protein 1) surface of ER and Lipid droplets, invo 37029364). In addition, AUP1 also co degradation of several key regulator 3-methylglutaryl-coenzyme A reduct Author Pu Milton To 27 Edmond Y Huang 29	vely, antigen vith citrate	ded proteins through autophagy (PMID: subiquitination and the subsequent e cholesterol biosynthetic enzyme 3-hydroxy Application WB	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




K-562 cells were subjected to SDS PAGE followed by western blot with 13726-1-AP (AUP1 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer using 13726-1-AP (AUP1 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver cancer using 13726-1-AP (AUP1 antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-AUP1 (IP:13726-1-AP, 3ug; Detection:13726-1-AP 1:500) with HEK-293 cells lysate 880ug.



Immunofluorescent analysis of A431 cells using 13726-1-AP (AUP1 antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.