

For Research Use Only

UBP1 Polyclonal antibody

Catalog Number: 13856-1-AP



Basic Information

Catalog Number: 13856-1-AP	GenBank Accession Number: BC047235	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 7342	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500
Source: Rabbit	Full Name: upstream binding protein 1 (LBP-1a)	
Isotype: IgG	Calculated MW: 60 kDa	
Immunogen Catalog Number: AG4830	Observed MW: 68 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A549 cells, human kidney tissue, HeLa cells, HepG2 cells, Jurkat cells, MCF-7 cells

IHC : human breast cancer tissue,

Background Information

UBP1, also known as Transcription factor LBP-1, is a 540 amino acid protein, which is expressed in adrenal tissue, JEG-3, NCI-H295A, Hep-G2 and HeLa cell lines. UBP1 localizes in nuclear and can be induced by HIV-1 infection of lymphocytes. UBP1 functions as a transcriptional activator in a promoter context-dependent manner and modulates the placental expression of CYP11A1. UBP1 is involved in regulation of the alpha-globin gene in erythroid cells and represses HIV-1 transcription by inhibiting the binding of TFIID to the TATA-box.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13856-1-AP (UBP1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 13856-1-AP (UBP1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).