

For Research Use Only

Siglec-7/CD328 Polyclonal antibody

Catalog Number: 13939-1-AP

2 Publications



Basic Information

Catalog Number:

13939-1-AP

Size:

150ul, Concentration: 450 ug/ml by Nanodrop and 287 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4220

GenBank Accession Number:

BC028150

GeneID (NCBI):

27036

UNIPROT ID:

Q9Y286

Full Name:

sialic acid binding Ig-like lectin 7

Calculated MW:

467 aa, 51 kDa

Observed MW:

65-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human placenta tissue, human liver tissue, U-937 cells, mouse placenta tissue

IHC: human tonsillitis tissue, human liver tissue, human placenta tissue

Background Information

Sialic acid binding Ig-like lectin 7 (Siglec-7), also known as CD328 or p75/AIRM-1, is a member of the Siglec family of glycan-recognition proteins. Siglec-7 is a type-I transmembrane protein consisting of three extracellular immunoglobulin-like domains that comprise an N-terminal V-set domain and two C2-set domains, a transmembrane region and a cytoplasmic tail containing two tyrosine residues embodied in immunoreceptor tyrosine-based inhibition motif-like motifs (PMID: 32322597; 10567377). It is mainly expressed on immune cells, with low levels on granulocytes, intermediate levels on monocytes, and relatively high levels on a major subset of natural killer cells and a minor subset of CD8+ T cells (PMID: 10567377). Siglec-7 is an inhibitory receptor that negatively regulates the function of NK cells and modulates the immune response through the interaction of sialic acid-containing ligands (PMID: 27312286).

Notable Publications

Author	Pubmed ID	Journal	Application
Kensuke Yamada	33240416	Oncol Lett	WB,IHC,IF
Rebecca Garnham	38448753	Commun Biol	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13939-1-AP (Siglec-7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis slide using 13939-1-AP (Siglec-7 Antibody) at dilution of 1:50.



Immunohistochemical analysis of paraffin-embedded human tonsillitis slide using 13939-1-AP (Siglec-7 Antibody) at dilution of 1:50.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 13939-1-AP (Siglec-7/CD328 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 13939-1-AP (Siglec-7/CD328 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).