For Research Use Only

ACSL1 Polyclonal antibody

Catalog Number: 13989-1-AP

Featured Product

57 Publications



Basic Information

Catalog Number: GenBank Accession Number:

 13989-1-AP
 BC050073

 Size:
 Genel D (NCBI):

 150ul , Concentration: 1000 ug/ml by 2180

Nanodrop; UNIPROT ID:
Source: P33121
Rabbit Full Name:

Isotype: acyl-CoA synthetase long-chain

IgG family member 1
Immunogen Catalog Number: Calculated MW:
AG5059 78 kDa

Observed MW: 68 kDa, 78 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:200-1:800 IF/ICC 1:400-1:1600

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: LO2 cells, mouse cerebellum tissue, mouse liver

tissue, rat liver tissue

IP: mouse liver tissue.

IHC: human lung squamous cell carcinoma tissue, human heart tissue, human liver cancer tissue

IF/ICC: HepG2 cells,

Background Information

ACSL1(Long-chain-fatty-acid--CoA ligase 1) is also named as FACL1, FACL2, LACS, LACS1, LACS2 and belongs to the ATP-dependent AMP-binding enzyme family. ACSL1 is a 75 kDa protein that is associated peripherally with the plasma membrane(Brian M Wiczer, etc., 2006). ACSL1 is abundantly expressed in tissues, such as liver and brown fat, that metabolize substantial amounts of triglycerides as fuel, and as such, a deficiency in ACSL1 function could have a more profound affect in those cells, resulting in hepatosteatosis and potentially increased very low density lipoprotein production by the liver or decreased thermogenic capacity in brown adipose tissue(PMID:19429676). An anti-rat ACSL1 antibody recognized a band of the predicted 68 kDa in high-speed supernatant from rat liver and in human and murine SMCs, monocyte-derived macrophages, and murine peritoneal macrophages (PMID:17259370). It has 2 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuxiang Sun	31590050	Colloids Surf B Biointerfaces	WB
De Huang	25242319	Cell Rep	WB
Qixue Wang	30279734	Theranostics	IF

Storage

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13989-1-AP (ACSL1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



IP result of anti-ACSL1 (IP:13989-1-AP, 4ug; Detection:13989-1-AP 1:1000) with mouse liver tissue lysate 3040 ug.



Immunohistochemical analysis of paraffinembedded human lung squamous cell carcinoma tissue slide using 13989-1-AP (ACSL1 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ACSL1 antibody (13989-1-AP) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).