For Research Use Only

p38 MAPK Polyclonal antibody

Catalog Number: 14064-1-AP

Featured Product

463 Publications



Basic Information

Catalog Number: GenBank Accession Number:

14064-1-APBC031574Size:GeneID (NCBI):150ul , Concentration: 600 ug/ml by1432

Nanodrop; UNIPROT ID:
Source: Q16539
Rabbit Full Name:

Isotype: mitogen-activated protein kinase 14

IgGCalculated MW:Immunogen Catalog Number:360 aa, 41 kDaAG5115Observed MW:

38-42 kDa

Applications

Tested Applications: WB, IHC, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, pig, rabbit, canine, chicken, goat,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, mouse heart tissue, Jurkat cells, RAW 264.7 cells, Neuro-2a cells, NIH/3T3 cells, rat spleen tissue, rat heart tissue, K-562 cells

Purification Method:

WB 1:2000-1:12000

IHC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

IHC: human colon cancer tissue, human liver cancer tissue

Background Information

MAPK14(mitogen-activated protein kinase 14) is also named as SAPK2A, p38MAPK, CSBP1, RK, p38, EXIP, Mxi2, CSBP2, PRKM14, PRKM15, CSPB1, p38ALPHA and belongs to the MAP kinase subfamily. MAPK14-signaling is a central pathway for the integration of instructive signals in dendritic cells for T(H)17 differentiation and inflammation(PMID:22231518). It plays an important role in the regulation of hematopoietic stem cellself-renewal in vitro and inhibition of MAPK14 activation with a small molecule inhibitor may represent a novel approach to promote ex vivo expansion of hematopoietic stem cell(PMID:21198398). This protein has some isoforms with MW 29-31 kDa, 35 kDa and 41 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Zemin Zhu	36175845	BMC Mol Cell Biol	WB
Xin-Sen Chen	36182039	Pharmacol Res	WB
Liping Wang	34559939	IUBMB Life	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




WB result of p38 antibody (14064-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-p38 transfected Jurkat cells.




Various lysates were subjected to SDS PAGE followed by western blot with 14064-1-AP (p38 MAPK antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 14064-1-AP (p38 MAPK antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 14064-1-AP (p38 MAPK antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 14064-1-AP (p38 MAPK antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 14064-1-AP (p38 MAPK antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).