

For Research Use Only

Frataxin Polyclonal antibody

Catalog Number: 14147-1-AP

Featured Product

20 Publications



Basic Information

Catalog Number:

14147-1-AP

Size:

150ul, Concentration: 500 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5316

GenBank Accession Number:

BC048097

GeneID (NCBI):

2395

UNIPROT ID:

Q16595

Full Name:

frataxin

Calculated MW:

23 kDa

Observed MW:

14 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IHC 1:300-1:1200

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF, CoIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : HCT 116 cells, Jurkat cells, K-562 cells, Raji cells, mouse liver tissue

IHC : human stomach tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Frataxin (FXN), also named as FRDA, X25, m81-FXN, d-FXN, m78-FXN and i-FXN, belongs to the frataxin family. It promotes the biosynthesis of heme and assembly and repair of iron-sulfur clusters by delivering Fe²⁺ to proteins involved in these pathways. FXN may play a role in the protection against iron-catalyzed oxidative stress through its ability to catalyze the oxidation of Fe²⁺ to Fe³⁺; the oligomeric form but not the monomeric form has in vitro ferroxidase activity. FXN is cleaved to be 4 chains. FXN is expressed in the cytoplasm as a 210 amino acid (AA) precursor protein (pFXN; 23 KDa) that is translocated into mitochondria where it is processed by two consecutive steps into iFXN (FXN 42-210; 19 KDa) and finally mFXN (81-210; 14.2 KDa), which is functional. (PMID: 26704351, PMID: 26671574)

Notable Publications

Author	Pubmed ID	Journal	Application
Jingjing Liu	33017621	Toxicology	WB
Daniel Fil	32586831	Dis Model Mech	WB
Yanjie Li	35708503	Hum Mol Genet	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1%BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14147-1-AP (Frataxin antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human stomach tissue slide using 14147-1-AP (Frataxin antibody) at dilution of 1:600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).