For Research Use Only

## NR2E3 Polyclonal antibody Catalog Number: 14246-1-AP Featured Product

Featured Product 9 Publications




| <b>Basic Information</b> | Catalog Number:<br>14246-1-AP  | GenBank Accession Number:<br>BC041421   | Purification Method:<br>Antigen affinity purification   |  |
|--------------------------|--|---|---|--|
|                          | Size:<br>150ul, Concentration: 600 ug/ml by<br>Nanodrop;<br>Source:<br>Rabbit<br>Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG5503  | GeneID (NCBI):<br>10002<br>UNIPROT ID:<br>Q9Y5X4<br>Full Name:<br>nuclear receptor subfamily 2, gro<br>member 3<br>Calculated MW:<br>45 kDa<br>Observed MW:<br>43-45 kDa  | Recommended Dilutions:<br>WB 1:2000-1:12000<br>IHC 1:20-1:200<br>IF/ICC 1:20-1:100  |  |
| Applications             | Tested Applications:<br>WB, IHC, IF/ICC, ELISA   | Positive Controls:  |   |  |
|                          | Cited Applications:  |   | WB : mouse retina tissue, Y79 cells, HepG2 cells, rat<br>retina tissue<br>IHC : human prostate cancer tissue,   |  |
|                          | WB, IHC  | IHC : hum   |   |  |
|                          | Species Specificity:   IF/ICC : photoreceptor progenitor of the photoreceptor photorecep |   |   |  |
|                          | Cited Species:<br>human, mouse, zebrafish<br>Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0  |   |   |  |
|                          |  |   |   |  |
| Background Information   | buffer pH 6.0<br>NR2E3, also known as PNR, encodes a<br>protein is part of a large family of nu<br>influences the development of photo<br>ranscriptional factor that is an activa  | clear receptor transcription factors<br>receptors and their differentiation<br>tor of rod development and repress<br>er of rod- and cone-specific genes,  | • • •   |  |
| Background Information   | buffer pH 6.0<br>NR2E3, also known as PNR, encodes a<br>protein is part of a large family of nu<br>influences the development of photo<br>ranscriptional factor that is an activa<br>binds the promoter region of a numb<br>specific phosphodiesterase beta sub  | clear receptor transcription factors<br>receptors and their differentiation<br>tor of rod development and repress<br>er of rod- and cone-specific genes,  | involved in signaling pathways. NR2E3   |  |
|                          | buffer pH 6.0<br>NR2E3, also known as PNR, encodes a<br>protein is part of a large family of nu<br>influences the development of photo<br>ranscriptional factor that is an activa<br>binds the promoter region of a numb<br>specific phosphodiesterase beta sub  | clear receptor transcription factors<br>receptors and their differentiation<br>tor of rod development and repress<br>er of rod- and cone-specific genes,<br>unit. [PMID:15689355]   | involved in signaling pathways. NR2E3<br>into rod and cone types, and acts as a<br>sor of cone development [PMID:20725840].<br>including rhodopsin, M- and S-opsin and ro                             |  |
|                          | buffer pH 6.0   NR2E3, also known as PNR, encodes a protein is part of a large family of nuinfluences the development of photor ranscriptional factor that is an activa binds the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of a numb specific phosphodiesterase beta subtractional factor of the promoter region of the promoter regio                          | clear receptor transcription factors<br>receptors and their differentiation<br>tor of rod development and repress<br>er of rod- and cone-specific genes,<br>unit. [PMID:15689355]   | involved in signaling pathways. NR2E3<br>into rod and cone types, and acts as a<br>sor of cone development [PMID:20725840].<br>including rhodopsin, M- and S-opsin and roc<br>Application             |  |
|                          | buffer pH 6.0   NR2E3, also known as PNR, encodes a protein is part of a large family of nuinfluences the development of photorranscriptional factor that is an activa binds the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the phosphodiesterase beta subtractional factor that is an activation of the phosphodiesterase beta subtractional factor that is an activation of the phosphodiesterase beta subtractional factor that is an activation of the phosphodiesterase beta subtractindef subtractivatindef subtractivation of the phosphosphodi subtra                          | clear receptor transcription factors<br>receptors and their differentiation<br>tor of rod development and repress<br>er of rod- and cone-specific genes,<br>unit. [PMID:15689355]<br>Demed ID Journal<br>378246 Sci Rep   | involved in signaling pathways. NR2E3<br>into rod and cone types, and acts as a<br>sor of cone development [PMID:20725840].<br>including rhodopsin, M- and S-opsin and ro<br>Application<br>WB        |  |
|                          | buffer pH 6.0   NR2E3, also known as PNR, encodes a protein is part of a large family of nuinfluences the development of photorranscriptional factor that is an activa binds the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the promoter region of a numb specific phosphodiesterase beta subtractional factor that is an activation of the phosphodiesterase beta subtractional factor that is an activation of the phosphodiesterase beta subtractional factor that is an activation of the phosphodiesterase beta subtractional factor that is an activation of the phosphodiesterase beta subtractindef subtractivatindef subtractivation of the phosphosphodi subtra                          | clear receptor transcription factors<br>receptors and their differentiation<br>tor of rod development and repress<br>er of rod- and cone-specific genes,<br>unit. [PMID:15689355]<br>omed ID Journal<br>378246 Sci Rep<br>318861 PLoS Genet<br>403063 J Biol Chem | involved in signaling pathways. NR2E3<br>into rod and cone types, and acts as a<br>sor of cone development [PMID:20725840].<br>including rhodopsin, M- and S-opsin and ro<br>Application<br>WB<br>IHC |  |


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




IF result of NR2E3 antibody (14246-1-AP, 1:50) with human photoreceptor progenitors cells.



Various lysates were subjected to SDS PAGE followed by western blot with 14246-1-AP (NR2E3 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer using 14246-1-AP (NR2E3 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human prostate cancer using 14246-1-AP (NR2E3 antibody) at dilution of 1:100 (under 40x lens).