For Research Use Only

PGT Polyclonal antibody Catalog Number: 14327-1-AP 3 Publications





14327-1-AP	BC041140	Antigen affinity purifi	cation
Size:	GenelD (NCBI):	Recommended Dilution	ons:
		WB 1:500-1:1000	
Nanodrop and 220 ug/mL by Bradford method using BSA as the standard;	UNIPROT ID: Q92959	IHC 1:50-1:500	IHC 1:50-1:500
Source:	Rabbit solute carrier organic anion		
	Calculated MW:		
5			
0 0	• • • • • •		
	65 kDa		
Tested Applications:	Positive Controls: WB : mouse heart tissue, NIH/3T3 cells, mouse liver		
			ls, mouse liver
WBIHC		-	
Species Specificity: human, mouse			
Cited Species: human, mouse, pig			
TE buffer pH 9.0; (*) Alternati	ively, antigen		
carrier that mediates either the release intracellular oxidation. Northern blo	ase of newly-synthesized PC t analysis revealed that PGT	s by cells and/or the uptake of PG mRNA has broad tissue distribution	s prior to on, suggesting i
Author Pub	med ID Journal	ŀ	Application
Hui Huang 330	000396 Dig Dis Sci	I	нс
Zhikun Zhan 348	38489 Int Immuno	pharmacol V	VB
Heewon Seo 246	95625 Biol Reprod	I	НС
Storage: Store at -20°C. Stable for one year af Storage Buffer:	iter shipment. 0% glycerol pH 7.3.		
	150ul, Concentration: 220 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG5594 Tested Applications: WB, IHC, ELISA Cited Applications: WB, IHC Species Specificity: human, mouse Cited Species: human, mouse, pig Note-IHC: suggested antigen of TE buffer pH 9.0; (*) Alternations Veffer pH 6.0 Prostaglandin transporter (PGT), also carrier that mediates either the release intracellular oxidation. Northern blorole in the transport and/ or metabol detect the endogenous PGT protein. Author Pub Hui Huang 3300 Zhikun Zhan 348 Heewon Seo 246	150ul, Concentration: 220 ug/ml by Bradford method using BSA as the standard; 6578 Source: UNIPROT ID: 092959 Source: Full Name: Rabbit solute carrier organic anio transporter family, memb IgG Calculated MW: Immunogen Catalog Number: 70 kDa AG5594 Observed MW: 65 kDa 0 Tested Applications: tis WB, IHC, ELISA WB Species Specificity: Human, mouse Numan, mouse Cited Species: human, mouse, pig Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Prostaglandin transporter (PGT), also known as SLCO2A1, has be carrier that mediates either the release of newly-synthesized PC intracellular oxidation. Northern blot analysis revealed that PCT role in the transport and/ or metabolic clearance of PG in diverse detect the endogenous PGT protein. Author Pubmed ID Journal Hui Huang 33000396 Dig Dis Sci Zhikun Zhan 34838489 Int Immuno Heewon Seo 24695625 Biol Reprod	150ul, Concentration: 220 ug/ml by Bradford method using BSA as the standard; 6578 WB 1:500-1:100 Manodrop and 220 ug/ml by Bradford method using BSA as the standard; Q32959 IHC 1:50-1:500 Source: Full Name: Q32959 Source: Full Name: Soute carrier organic anion Isotype: transporter family, member 2A1 IgG IgG Calculated MW: Diserved MW: AG5594 Observed MW: 65 kDa Tested Applications: Positive Controls: WB, IHC, EUSA WB: mouse heart tissue, NIH/3T3 cell Cited Applications: WB: mouse heart tissue, NIH/3T3 cell Cited Applications: WB: mouse heart tissue, NIH/3T3 cell WB, IHC IHC : human stomach tissue, Species Specificity: IHC : human stomach tissue, human, mouse, pig Note-IHC: suggested antigen retrieval with T Estudy and transporter (PCT), also known as SLCO2A1, has been identified as a functional prostic carrier that mediates either the release of newly-synthesized PGs by cells and/or the uptake of PG in intracellular oxidation. Northern blot analysis revealed that PGT mRNA has broad tissue distribution role in the transport and/or metabolic clearance of PG in diverse tissues (8/78/67/7). This antibody detect the endogenous PGT protein. Author Pu

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 14327-1-AP (PGT antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Various lysates were subjected to SDS PAGE followed by western blot with 14327-1-AP (PGT antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 14327-1-AP (PGT antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).