

For Research Use Only

PON2 Polyclonal antibody

Catalog Number: 14379-1-AP **4 Publications**



Basic Information

Catalog Number: 14379-1-AP	GenBank Accession Number: BC046160	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 200 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 5445	Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Source: Rabbit	Full Name: paraoxonase 2	IHC 1:20-1:200
Isotype: IgG	Calculated MW: 39 kDa	IF 1:20-1:200
Immunogen Catalog Number: AG5759	Observed MW: 39 kDa	

Applications

Tested Applications: FC, IF, IHC, IP, WB, ELISA	Positive Controls: WB : human liver tissue, L02 cells
Cited Applications: IF, IHC, WB	IP : L02 cells,
Species Specificity: human	IHC : human liver cancer tissue,
Cited Species: human, mouse	IF : HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

PON2 (Serum paraoxonase/arylesterase 2) has antioxidant activity and can prevent LDL lipid peroxidation, reverse the oxidation of mildly oxidized LDL, and inhibit the ability of MM-LDL to induce monocyte chemotaxis. Highest levels of PON2 protein are found in the mouse lung and small intestine, followed by the heart and liver, while lower levels are present in the testis, kidney and brain. PON2 expression in tissues from female mice is always significantly higher than in male animals. There are also some reports showing two bands of 43 kDa and 53 kDa to be detected through western blot as the two isoforms of this protein. (PMID:21354197).

Notable Publications

Author	Pubmed ID	Journal	Application
Henning Hagmann	36429053	Cells	IHC
Henning Hagmann	24421402	FASEB J	IHC
Xueqi Chen	37054540	Biomed Pharmacother	WB, IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human liver tissue were subjected to SDS PAGE followed by western blot with 14379-1-AP (PON2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of HepG2 cells, using PON2 antibody 14379-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).




IP Result of anti-PON2 (IP:14379-1-AP, 4ug; Detection:14379-1-AP 1:500) with L02 cells lysate 3200ug.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 14379-1-AP (PON2 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 14379-1-AP (PON2 Antibody) at dilution of 1:200 (under 40x lens).



1X10⁶ HepG2 cells were stained with 0.2ug PON2 antibody (14379-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.