For Research Use Only

HE4 Polyclonal antibody Catalog Number: 14406-1-AP 7 Publications





Basic Information	Catalog Number: 14406-1-AP	GenBank Accession Number: BC046106	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 500 ug/ml by	10406	IHC: 1:250-1:1000	
	Nanodrop and 207 ug/ml by Bradford method using BSA as the standard;	UNIFICITID.	IF/ICC: 1:50-1:500	
	Source:	Q14508		
	Rabbit	Full Name: WAP four-disulfide core domain 2		
	lsotype:	Calculated MW:		
	IgG Immunogen Catalog Number:	13 kDa		
	AG5684			
Applications	Tested Applications:	Positive Controls:		
	IHC, IF/ICC, ELISA IHC : hu		an ovary tumor tissue, human testis tissue	
	Cited Applications: WB, IHC	IF/ICC : MCF-7 cells,		
	Species Specificity: human			
	Cited Species:			
	human, mouse			
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen		
Background Information	TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0 WFDC2, also named as HE4, is a mem contains eight cysteines forming four in many family members. HE4 is exp	rely, antigen ith citrate aber of the WFDC domain family. T disulfide bonds at the core of the ressed in a number of normal tissu arian, colon, breast, lung and renal	protein, and functions as a protease inhibite es, and it is also highly expressed in a	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 WFDC2, also named as HE4, is a mem contains eight cysteines forming four in many family members. HE4 is exp number of tumors cells lines, such ov- to as a biomarker for ovarian carcinor	rely, antigen ith citrate aber of the WFDC domain family. T disulfide bonds at the core of the ressed in a number of normal tissu arian, colon, breast, lung and renal	protein, and functions as a protease inhibit es, and it is also highly expressed in a	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0WFDC2, also named as HE4, is a mem contains eight cysteines forming four in many family members. HE4 is exp number of tumors cells lines, such ov- to as a biomarker for ovarian carcinouAuthorPub	rely, antigen ith citrate ober of the WFDC domain family. T disulfide bonds at the core of the ressed in a number of normal tissu arian, colon, breast, lung and renal ma.	protein, and functions as a protease inhibit es, and it is also highly expressed in a cells lines. In some study, HE4 was referre	
Background Information	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0WFDC2, also named as HE4, is a mem contains eight cysteines forming four in many family members. HE4 is exp number of tumors cells lines, such ov- to as a biomarker for ovarian carcinomAuthorPub Ran ChenRan Chen292	wely, antigen ith citrate aber of the WFDC domain family. T disulfide bonds at the core of the ressed in a number of normal tissu arian, colon, breast, lung and renal ma. med ID Journal	cells lines. In some study, HE4 was referred Application WB	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0WFDC2, also named as HE4, is a mem contains eight cysteines forming four in many family members. HE4 is exp number of tumors cells lines, such ov- to as a biomarker for ovarian carcinorAuthorPub Ran Chen292 20andan Fang316	rely, antigen ith citrate aber of the WFDC domain family. T disulfide bonds at the core of the p ressed in a number of normal tissu arian, colon, breast, lung and renal ma. med ID Journal 28604 Oncotarget	protein, and functions as a protease inhibit es, and it is also highly expressed in a cells lines. In some study, HE4 was referre Application WB	
Notable Publications	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0WFDC2, also named as HE4, is a mem contains eight cysteines forming four in many family members. HE4 is exp number of tumors cells lines, such ov- to as a biomarker for ovarian carcinorAuthorPub Ran Chen292Dandan FangJian Fang352	weight antigen ith citrate aber of the WFDC domain family. T disulfide bonds at the core of the ressed in a number of normal tissu arian, colon, breast, lung and renal ma. med ID Journal 28604 Oncotarget 06687 Biosens Bioelectre	protein, and functions as a protease inhibit es, and it is also highly expressed in a cells lines. In some study, HE4 was referre Application WB	
Notable Publications	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 WFDC2, also named as HE4, is a mem contains eight cysteines forming four in many family members. HE4 is exp number of tumors cells lines, such over to as a biomarker for ovarian carcinot Author Pub Ran Chen 292 Dandan Fang 316 Jian Fang 352 Storage: Storage Buffer:	rely, antigen ith citrate aber of the WFDC domain family. T disulfide bonds at the core of the p ressed in a number of normal tissu arian, colon, breast, lung and renal ma. med ID Journal 28604 Oncotarget 06687 Biosens Bioelectro 97435 Food Funct	protein, and functions as a protease inhibit es, and it is also highly expressed in a cells lines. In some study, HE4 was referre Application WB	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 WFDC2, also named as HE4, is a mem contains eight cysteines forming four in many family members. HE4 is exp number of tumors cells lines, such ov. to as a biomarker for ovarian carcinor Author Pub Ran Chen 292 Dandan Fang 316 Jian Fang 352 Storage: Storage:	rely, antigen ith citrate aber of the WFDC domain family. T disulfide bonds at the core of the p ressed in a number of normal tissu arian, colon, breast, lung and renal ma. med ID Journal 28604 Oncotarget 06687 Biosens Bioelectro 97435 Food Funct er shipment.	protein, and functions as a protease inhibit es, and it is also highly expressed in a cells lines. In some study, HE4 was referre Application WB	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data







Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 14406-1-AP (HE4 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 14406-1-AP (HE4 antibody) at dilution of 1:200 and Coralite488-Conjugated Goat Anti-Rabbit IgG(H+L).