For Research Use Only

TFIP11 Polyclonal antibody Catalog Number:14436-1-AP





Basic Information	Catalog Number: 14436-1-AP	GenBank Accession Number: BC033080	Purification Method: Antigen affinity purification				
	Size: 150ul, Concentration: 200 ug/ml by Nanodrop and 160 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI):	Recommended Dilutions: WB 1:500-1:1000 IF/ICC 1:50-1:500				
		24144 UNIPROT ID: O9UBB9					
				Source:	Full Name:		
	Rabbit Isotype: IgG Immunogen Catalog Number: AG5821	tuftelin interacting protein 11 Calculated MW: 97 kDa Observed MW: 97 kDa					
				A 11 41	Tested Applications:	Desitive C	antrol c
				Applications	WB, IF/ICC, ELISA		Positive Controls: WB : HeLa cells, K-562 cells, MCF-7 cells, mouse testi: tissue
	Species Specificity: human, mouse, rat						
		IF/ICC : He	La cells,				
Background Information	acid protein, which contains one G-pa nucleus and cytoplasm. TFIP11 is inv	acting protein 11 or Septin and tufte tch domain and belongs to the TFP olved in pre-mRNA splicing, specifi ay play a role in the differentiation	La cells, Lin-interacting protein 1, is a 837 amino 11/STIP family. TFIP11 may localizes in th cally in spliceosome disassembly during of ameloblasts and odontoblasts or in the				
Background Information	acid protein, which contains one G-pa nucleus and cytoplasm. TFIP11 is inv late-stage splicing events. TFIP11 ma forming of the enamel extracellular r Storage: Store at -20°C. Stable for one year aft Storage Buffer:	acting protein 11 or Septin and tufte tch domain and belongs to the TFP olved in pre-mRNA splicing, specifi ay play a role in the differentiation matrix. er shipment.	lin-interacting protein 1, is a 837 amino 11/STIP family. TFIP11 may localizes in tr cally in spliceosome disassembly during				
	acid protein, which contains one G-pa nucleus and cytoplasm. TFIP11 is inv late-stage splicing events. TFIP11 ma forming of the enamel extracellular r Storage: Store at -20°C. Stable for one year aft	acting protein 11 or Septin and tufte tch domain and belongs to the TFP olved in pre-mRNA splicing, specifi yy play a role in the differentiation natrix. er shipment. % glycerol pH 7.3.	lin-interacting protein 1, is a 837 amino 11/STIP family. TFIP11 may localizes in th cally in spliceosome disassembly during				

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





HeLa cells were subjected to SDS PAGE followed by western blot with 14436-1-AP (TFIP11 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 14436-1-AP (TFIP11 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).