For Research Use Only

MSRA Polyclonal antibody

Catalog Number: 14547-1-AP 13 Publications



Basic Information

Catalog Number: 14547-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul , Concentration: 350 ug/ml by

BC054033

WB 1:5000-1:10000

Nanodrop and 227 ug/ml by Bradford $\,$ UNIPROT ID:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

method using BSA as the standard;

Q9UJ68

protein lysate

Source: Rabbit

Full Name:

IHC 1:50-1:500

IF/ICC 1:50-1:500

Isotype

methionine sulfoxide reductase A Calculated MW:

26 kDa

Immunogen Catalog Number: AG6053

Observed MW:

24-26 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications

WB, IHC, IF, IP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HT-1080 cells, human kidney tissue, mouse

kidney tissue, rat kidney tissue

IP: mouse kidney tissue, IHC: human colon tissue,

IF/ICC: HEK-293 cells,

Background Information

Methionine sulfoxide reductase A(MSRA), is a ubiquitous and highly conserved protein that carries out the enzymatic reduction of methionine sulfoxide to methionine.MSRA has some isoforms with MW of 19-26 kDa. MSRA functions in the repair of oxidatively damaged proteins to restore biological activity. Transfection of MSRA reduced colony formation and the invasiveness of HCC cell lines.

Notable Publications

Author	Pubmed ID	Journal	Application
Lo Lai	31606431	Free Radic Biol Med	WB
Qi Zhang	36233020	Int J Mol Sci	IF,WB
Dani Qin	36448374	Cell Biol Int	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using 14547-1-AP (MSRA antibody), at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).




Various lysates were subjected to SDS PAGE followed by western blot with 14547-1-AP (MSRA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-MSRA (IP:14547-1-AP, 4ug; Detection:14547-1-AP 1:600) with mouse kidney tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 14547-1-AP (MSRA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 14547-1-AP (MSRA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).