For Research Use Only

APOBEC3B Polyclonal antibody

Catalog Number:14559-1-AP 4 Publications




Basic Information	Catalog Number: 14559-1-AP	GenBank Accession Number: BC053859	Purification Method: Antigen affinity purification			
	Size: 150ul, Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI):	Recommended Dilutions:			
		y 9582 WB 1:1000-1:4000 UNIPROT ID: Q9UH17 Full Name: apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3B				
				Immunogen Catalog Number: AG6115	Calculated MW: 46 kDa	
					Observed MW: 46 kDa	
				Applications	Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, HepG2 cells
	Cited Applications: WB, IP, IHC					
	Species Specificity: human					
	Cited Species: human, mouse					
Background Information	human, mouse The DNA cytosine deaminase APOBE breast, head/neck, cervix, bladder, lu	ng, ovary, and to lesser extents e response to virus infection. Af	POBEC3B has 3 isoforms produced by alternativ			
	human, mouse The DNA cytosine deaminase APOBE breast, head/neck, cervix, bladder, lu effector protein in the innate immune splicing with the MW of 46, 29, 43 kDa	ng, ovary, and to lesser extents e response to virus infection. Af a. It can also exist as a dimer an	additional cancer types. This enzyme is also ar POBEC3B has 3 isoforms produced by alternative nd can interact with APOBEC3G.			
	human, mouse The DNA cytosine deaminase APOBE breast, head/neck, cervix, bladder, lu effector protein in the innate immune splicing with the MW of 46, 29, 43 kDa Author Pub	ng, ovary, and to lesser extents e response to virus infection. Af a. It can also exist as a dimer an med ID Journal	additional cancer types. This enzyme is also ar POBEC3B has 3 isoforms produced by alternative nd can interact with APOBEC3G. Application			
	human, mouse The DNA cytosine deaminase APOBE breast, head/neck, cervix, bladder, lu effector protein in the innate immune splicing with the MW of 46, 29, 43 kDa Author Pub Hao Zhang 338	ng, ovary, and to lesser extents e response to virus infection. Af a. It can also exist as a dimer an med ID Journal 42328 Front Oncol	additional cancer types. This enzyme is also ar POBEC3B has 3 isoforms produced by alternative and can interact with APOBEC3G. Application IHC			
Background Information Notable Publications	human, mouse The DNA cytosine deaminase APOBE breast, head/neck, cervix, bladder, lu effector protein in the innate immune splicing with the MW of 46, 29, 43 kDa Author Pub Hao Zhang 338 Qiong Fan 336	ng, ovary, and to lesser extents e response to virus infection. Af a. It can also exist as a dimer an med ID Journal	additional cancer types. This enzyme is also ar POBEC3B has 3 isoforms produced by alternative nd can interact with APOBEC3G. Application			
	human, mouse The DNA cytosine deaminase APOBE breast, head/neck, cervix, bladder, lu effector protein in the innate immune splicing with the MW of 46, 29, 43 kDa Author Pub Hao Zhang 338 Qiong Fan 336	ng, ovary, and to lesser extents e response to virus infection. Af a. It can also exist as a dimer an med ID Journal 42328 Front Oncol 31046 Mol Carcinog 62729 J Med Genet er shipment. % glycerol pH 7.3.	additional cancer types. This enzyme is also ar POBEC 3B has 3 isoforms produced by alternative and can interact with APOBEC 3G. Application IHC IHC, WB, IP			

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14559-1-AP (APOBEC3B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.