

For Research Use Only

GSTO2 Polyclonal antibody

Catalog Number: 14562-1-AP

8 Publications



Basic Information

Catalog Number:

14562-1-AP

Size:

150ul , Concentration: 600 ug/ml by Nanodrop and 367 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6120

GenBank Accession Number:

BC056918

GeneID (NCBI):

119391

UNIPROT ID:

Q9H4Y5

Full Name:

glutathione S-transferase omega 2

Calculated MW:

28 kDa

Observed MW:

28-31 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IF/ICC 1:20-1:200

Applications

Tested Applications:

WB, IF/ICC, ELISA

Cited Applications:

WB, ColP, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : HCT 116 cells, mouse testis tissue, rat testis tissue, mouse skeletal muscle tissue

IF/ICC : HeLa cells,

Notable Publications

Author	Pubmed ID	Journal	Application
Cheng-Niu Wang	32987309	Bioorg Chem	WB
Rojanathammanee Lalida L	24285747	J Gerontol A Biol Sci Med Sci	WB
Lalida Rojanathammanee	25001375	Age (Dordr)	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 14562-1-AP (GSTO2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 14562-1-AP (GSTO2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).

Various lysates were subjected to SDS PAGE followed by western blot with 14562-1-AP (GSTO2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.