For Research Use Only

AFF4 Polyclonal antibody

Catalog Number: 14662-1-AP

Featured Product

12 Publications



Basic Information

Catalog Number: GenBank Accession Number:

14662-1-AP BC063007 GeneID (NCBI):

150ul , Concentration: 1100 $\mu g/ml$ by 27125

Nanodrop;

AF4/FMR2 family, member 4 Source:

Rabbit Calculated MW: Isotype: 127 kDa IgG Observed MW: Immunogen Catalog Number: 140-150 kDa

AG6336

Applications Tested Applications:

IP, WB, ELISA

Cited Applications: IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:5000-1:50000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Positive Controls:

WB: HEK-293 cells, HeLa cells, HepG2 cells, mouse

brain tissue, rat brain tissue

IP: HeLa cells,

Background Information

Transcriptional controled by RNA polymerase II (Pol II) is a multi-step process requiring the concerted action of multiple factors and contacts with the DNA template for the proper synthesis of nascent RNA [PMID:12676794]. ELL1 was demonstrated to be a Pol II elongation factor capable of enhancing the catalytic rate of transcription elongation by reducing transient pausing by the enzyme [PMID:8596958]. AFF4 was initially identified as a fusion partner of the MLL (mixed-lineage leukemia) gene participated in infant acute lymphoblastic leukemia, and a component of the ELL/p-TEFb elongation complex. AFF4 interacts with positive transcription elongation factor-b while also repressing Tat transactivation of HIV-1 [PMID:17389929].

Notable Publications

Author	Pubmed ID	Journal	Application
Jun Wu	28878233	Sci Rep	WB
Bin Zheng	33446572	Genes Dev	WB
Dan Tang	32128251	Cell Discov	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14662-1-AP (AFF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



IP result of anti-AFF4(IP:14662-1-AP, 4ug; Detection:14662-1-AP 1:20000) with HeLa cells lysate 1880 ug.