

For Research Use Only

RBM42 Polyclonal antibody

Catalog Number: 15407-1-AP



Basic Information

Catalog Number:

15407-1-AP

Size:

150ul , Concentration: 550 ug/ml by Nanodrop and 380 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG7667

GenBank Accession Number:

BC004204

GeneID (NCBI):

79171

UNIPROT ID:

Q9BTD8

Full Name:

RNA binding motif protein 42

Calculated MW:

50 kDa

Observed MW:

65-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human, mouse, rat

Positive Controls:

WB : HepG2 cells, HEK-293 cells, HeLa cells, human liver tissue, NIH/3T3 cells

Background Information

RBM42, also known as RNA-binding motif protein 42, is a member of the RNA-binding protein family that plays critical roles in post-transcriptional regulation of gene expression. Emerging studies highlight RBM42's involvement in pre-mRNA splicing through its association with the spliceosome complex, where it contributes to the precise removal of introns and assembly of mature mRNA. Beyond splicing, RBM42 has been implicated in cellular stress responses, cell cycle regulation, and apoptosis. Dysregulation of RBM42 is linked to developmental disorders and cancers; for instance, mutations in RBM42 are associated with neurodevelopmental abnormalities, while its overexpression or loss correlates with tumor progression in certain malignancies.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 15407-1-AP (RBM42 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.