For Research Use Only

CYFIP1/2 Polyclonal antibody

Catalog Number:16011-1-AP 2 Publications




	16011-1-AP	BC011762		Antigen affinity purification	
	Size:	GenelD (NCBI): 26999		Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:300-1:1200	
	150ul , Concentration: 350 ug/ml by				
	Nanodrop and 193 ug/ml by Bradford method using BSA as the standard;				
					Source:
	Rabbit				cytoplasmic FMR1 interacting protein 2
	Isotype:				
	IgG Immunogen Catalog Number: AG8749	Calculated M	N:		
		148 kDa			
		Observed MW 130 kDa	1		
	Applications	Tested Applications:	Positive Controls:		
WB, IHC, FC (Intra), IP, ELISA Cited Applications:			WB : human tissue	: human brain tissue, mouse brain tissue, rat brain sue mouse brain tissue,	
WB			IP : mouse b		
Species Specificity:			IHC : mouse	e brain tissue, human gliomas tissue,	
		human, mouse, rat human ki		Iney tissue	
Cited Species: human					
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	CYFIP1/2 belongs to the CYFIP family and is involved in T-cell adhesion and p53-dependent induction of apoptosis CYFIP1 contains 1,253 amino acids and shares 88% sequence identity with CYFIP2. CYFIP1 and CYFIP2 are members of a highly conserved protein family and share approximately 99% sequence identity with their mouse orthologs. CYFIP1 has been shown to interact with the small GTPase RAC1, which is implicated in development and maintenance of neuronal structures. Consistent with FMRP and RAC1 localization in dendritic fine structures, CYFIP1/2 are present in synaptosomal extracts. This antibody can recognize both CYFIP1 and CYFIP2.				
Notable Publications	Author Pub	omed ID	Journal	Application	
	Nachmany Henny H 224	430187	Dis Markers	WB	
	Mirjana Malnar Črnigoj 377	717009	Nat Commun	WB	
Storage	Storage: Store at -20°C. Stable for one year after Storage Buffer:				
Juliage	PBS with 0.02% sodium azide and 50°	% glycerol pH 1	1.3.		
Joinage	PBS with 0.02% sodium azide and 50 Aliquoting is unnecessary for -20°C st		/.3.		


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human brain tissue were subjected to SDS PAGE followed by western blot with 16011-1-AP (CYFIP2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 16011-1-AP (CYFIP1/2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16011-1-AP (CYFIP1/2 antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-CYFIP1/2 (IP:16011-1-AP, 4ug; Detection:16011-1-AP 1:700) with mouse brain tissue lysate 4000ug.



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug Anti-Human CYFIP1/2 (16011-1-AP) and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).