For Research Use Only

GALNTL2 Polyclonal antibody

Catalog Number: 16900-1-AP



Basic Information

Catalog Number: GenBank Accession Number:

16900-1-AP Antigen affinity purification BC014789 Size: GeneID (NCBI): Recommended Dilutions: WB 1:500-1:1000 150ul , Concentration: 133 μ g/ml by 117248 IHC 1:50-1:500

Nanodrop and 133 µg/ml by Bradford Full Name: method using BSA as the standard;

UDP-N-acetyl-alpha-Dgalactosamine:polypeptide N-Rabbit acetylgalactosaminyltransferase-like

Isotype:

IgG Calculated MW: 639 aa, 73 kDa Immunogen Catalog Number: Observed MW: AG10434 65-70 kDa

Applications

Tested Applications:

IHC, WB, ELISA Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, human liver tissue

IHC: mouse small intestine tissue, human kidney tissue, human testis tissue, human placenta tissue, human brain tissue, human lung tissue, human ovary

Purification Method:

Background Information

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 16900-1-AP (GALNTL2 antibody) at dilution of 1:200 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 16900-1-AP (GALNTL2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse small intestine tissue slide using 16900-1-AP (GALNTL2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse small intestine tissue slide using 16900-1-AP (GALNTL2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 16900-1-AP (GALNTL2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).