For Research Use Only

CD11c Polyclonal antibody Catalog Number: 17342-1-AP 69 Publications




Basic Information	Catalog Number: 17342-1-AP	BC038237 // GeneID (NCBI): // 3687 //		Purification Method: Antigen affinity purification				
Basic Information								
Applications	Tested Applications: WB, IHC, ELISA	Positive Controls: WB : THP-1 cells, human plasma IHC : human tonsillitis tissue,						
	Cited Applications: WB, IHC, IF							
	Species Specificity: human Cited Species: human, pig Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits CD11c/Integrin alpha X is a 145-150 kDa type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, NK cells and dendritic cells. Integrin alpha X/beta 2 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis.							
					Background Information			
					Notable Publications	Author Pul	bmed ID	Journal
Hao Zhang 360						093059	iScience	IF
Haoran Wang 350	087966	Bioact Mater						
Ryutaro Hara 33	178566	Asia Pac Allergy	IHC,IF					
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol, pH7.3							
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	storage						
For technical support and original validation dat	ta for this product please contact:		This product is	exclusively available under Proteintec				

W: ptglab.com


in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 17342-1-AP (CD11c/Integrin Alpha X antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 17342-1-AP (CD11c/Integrin Alpha X antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

THP-1 cells were subjected to SDS PAGE followed by western blot with 17342-1-AP (CD11c/Integrin Alpha X antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.