For Research Use Only

IL-12/IL-23 p40 Polyclonal antibody

Catalog Number: 17645-1-AP 4 Publications





Basic Information	Catalog Number: 17645-1-AP	GenBank Accession Number: BC067498	Purification Method: Antigen affinity purification			
	Size:	GenelD (NCBI):	Recommended Dilutions:			
	150ul, Concentration: 400 ug/ml by Nanodrop and 293 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number:	3593	WB 1:1000-1:4000			
		UNIPROT ID: P29460				
		Full Name:				
		interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40) Calculated MW:				
				AG11837	328 aa, 37 kDa	
					Observed MW: 39 kDa	
	Applications	Tested Applications:	Positive Controls:			
		WB, FC (Intra), ELISA	WB : Recombinant protein,			
Cited Applications: WB, IF						
Species Specificity: human						
Cited Species:						
	human, mouse					
Background Information	human, mouse IL-12 and IL-23 are heterodimeric cytr of the IL-12 p40 subunit linked to the 12R), which comprises the IL-12Rβ1 a	IL-12 p35 subunit, and the heterodi nd IL-12Rβ2 subunits. IL-23 is comp hals through IL-23R and IL-12Rβ1 (PI modimer which can act as a potent ed by antigen-presenting cells, such	mer signals through the IL-12 receptor (IL- osed of the IL-23 p19 subunit and the IL-12 MID: 11114383; 26121196). IL-12/IL-23 p44 IL-12 antagonist (PMID: 8958912; a s dendritic cells (DCs), monocytes,			
	human, mouse IL-12 and IL-23 are heterodimeric cyte of the IL-12 p40 subunit linked to the 12R), which comprises the IL-12Rβ1 a p40 (IL-12/23p40) subunit, which sign also exists as a monomer and as a ho 18783467). IL-12/IL-23 p40 is produce macrophages, neutrophils and, to a le	IL-12 p35 subunit, and the heterodi nd IL-12Rβ2 subunits. IL-23 is comp hals through IL-23R and IL-12Rβ1 (PI modimer which can act as a potent ed by antigen-presenting cells, such esser extent, B cells (PMID: 2047691	mer signals through the IL-12 receptor (IL- osed of the IL-23 p19 subunit and the IL-12 MID: 11114383; 26121196). IL-12/IL-23 p44 IL-12 antagonist (PMID: 8958912; as dendritic cells (DCs), monocytes, 8).			
	human, mouse IL-12 and IL-23 are heterodimeric cytr of the IL-12 p40 subunit linked to the 12R), which comprises the IL-12Rβ1 a p40 (IL-12/23p40) subunit, which sign also exists as a monomer and as a ho 18783467). IL-12/IL-23 p40 is produce macrophages, neutrophils and, to a le	IL-12 p35 subunit, and the heterodi nd IL-12Rβ2 subunits. IL-23 is comp hals through IL-23R and IL-12Rβ1 (PI modimer which can act as a potent ed by antigen-presenting cells, such esser extent, B cells (PMID: 2047691 med ID Journal	mer signals through the IL-12 receptor (IL- osed of the IL-23 p19 subunit and the IL-12 MID: 11114383; 26121196). IL-12/IL-23 p4 IL-12 antagonist (PMID: 8958912; a as dendritic cells (DCs), monocytes, 8). Application			
Background Information	human, mouse IL-12 and IL-23 are heterodimeric cyte of the IL-12 p40 subunit linked to the 12R), which comprises the IL-12Rβ1 a p40 (IL-12/23p40) subunit, which sign also exists as a monomer and as a ho 18783467). IL-12/IL-23 p40 is produce macrophages, neutrophils and, to a le Author Pub Xinlong Zang 333	IL-12 p35 subunit, and the heterodi nd IL-12Rβ2 subunits. IL-23 is comp hals through IL-23R and IL-12Rβ1 (PI modimer which can act as a potent ed by antigen-presenting cells, such esser extent, B cells (PMID: 2047691	osed of the IL-23 p19 subunit and the IL-12 MID: 11114383; 26121196). IL-12/IL-23 p40 IL-12 antagonist (PMID: 8958912; a as dendritic cells (DCs), monocytes, 8). Application IF			
	human, mouse	IL-12 p35 subunit, and the heterodi nd IL-12Rβ2 subunits. IL-23 is comp hals through IL-23R and IL-12Rβ1 (PI modimer which can act as a potent ed by antigen-presenting cells, such asser extent, B cells (PMID: 2047691 med ID Journal 38567 Int J Pharm	mer signals through the IL-12 receptor (IL- osed of the IL-23 p19 subunit and the IL-12 MID: 11114383; 26121196). IL-12/IL-23 p4 IL-12 antagonist (PMID: 8958912; n as dendritic cells (DCs), monocytes, 8). Application IF			

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Recombinant protein IL-12B (Eg0431), IL-12 (Eg0575), IL-12A (Ag17625) were subjected to SDS PAGE followed by western blot with 17645-1-AP (IL-12B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours 1x10^6 Jurkat cells were intracellularly stained with 0.4 ug IL-12/IL-23 p40 Polyclonal antibody (17645-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).