

ICAM3/CD50 Polyclonal antibody

Catalog Number: 17699-1-AP

Basic Information

Catalog Number:

17699-1-AP

Size:

150ul, Concentration: 350 ug/ml by Nanodrop and 193 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG11955

GenBank Accession Number:

BC058903

GeneID (NCBI):

3385

UNIPROT ID:

P32942

Full Name:

intercellular adhesion molecule 3

Calculated MW:

547 aa, 60 kDa

Observed MW:

110-160 kDa, 60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2400

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

Applications

Tested Applications:

WB, IHC, IP, ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, Ramos cells, U-937 cells

IP: Jurkat cells,

IHC: human colon tissue,

Background Information

ICAM3 (Intercellular adhesion molecule 3) is also named as CD50, CDw50 and ICAM-R. ICAM3 is a member of the intercellular adhesion molecule (ICAM) family. ICAM3, being expressed constitutively on T cells and other leukocytes, is a potent signalling molecule and a major ligand in the initiation of T cell-mediated immune responses (PMID: 1448173, PMID: 11812993). ICAM3 is the natural ligand for both DC-SIGN and L-SIGN (PMID: 17570115). ICAM3 is abundantly and constitutively expressed in all leukocytes and is probably the most important ligand for LFA-1 in initiating immune responses, and ICAM3 is a typical type I transmembrane glycoprotein (PMID: 39991572). The promoter region of ICAM3 can be bound by co-transcription factors such as ETs and C/EBP, promoting macrophage maturation as well as monocyte extravasation (PMID: 22479382). ICAM3 is widely expressed not only in lymphocytes such as T and B cells but also in many endothelial cells and dendritic cells (PMID: 39991572).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 17699-1-AP (ICAM3/CD50 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon using 17699-1-AP (ICAM3/CD50 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human colon using 17699-1-AP (ICAM3 antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-ICAM3/CD50 (IP:17699-1-AP, 4ug; Detection:17699-1-AP 1:500) with Jurkat cells lysate 4000 ug.