For Research Use Only

DDX10 Polyclonal antibody

Catalog Number: 17857-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number: GenBank Accession Number: 17857-1-AP BC091521

Size: GeneID (NCBI):

150ul , Concentration: 550 µg/ml by 1662

Nanodrop and 273 µg/ml by Bradford method using BSA as the standard; DEAD (Asp.

nethod using BSA as the standard; DEAD (Asp-Glu-Ala-Asp) box

Source: polypeptide 10
Rabbit Calculated MW:
Isotype: 875 aa, 101 kDa
IgG Observed MW:

Immunogen Catalog Number: 101 kDa

AG12317

Positive Controls:

WB: SMMC-7721 cells, HeLa cells

 $\hbox{IHC:} \ mouse \ test is \ tissue, human \ kidney \ tissue$

Purification Method:

WB 1:200-1:1000

IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

IHC, WB

Species Specificity: human, mouse, rat Cited Species:

human

Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate buffer pH 6.0

Background Information

DDX10, DEAD-box helicase 10, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are involved in cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. Fusion of this gene and the nucleoporin gene NUP98 is found in leukemogenesis (PMID:20339440).

Notable Publications

Author	Pubmed ID	Journal	Application
Xin Zhou	35109823	Cancer Cell Int	WB,IHC
Muhuizi Gai	26713367	Biochem Biophys Res Commun	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




SMMC-7721 cells were subjected to SDS PAGE followed by western blot with 17857-1-AP (DDX10 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 17857-1-AP (DDX10 Antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 17857-1-AP (DDX10 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 17857-1-AP (DDX10 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).