### For Research Use Only

# Galectin-9 Polyclonal antibody

Catalog Number: 17938-1-AP

Featured Product

10 Publications



**Purification Method:** 

**Basic Information** 

Catalog Number: GenBank Accession Number:

17938-1-AP BC110340 Antigen affinity purification
Size: GeneID (NCBI): Recommended Dilutions:

150ul , Concentration: 300 µg/ml by 3965 WB 1:500-1:1000

Nanodrop; Full Name: IP 0.5-4.0 ug for IP and 1:500-1:3000

Source: lectin, galactoside-binding, soluble, 9 for WB
Rabbit Calculated MW:

Isotype:323 aa, 36 kDaIgGObserved MW:Immunogen Catalog Number:45 kDa

AG12209

**Applications** 

Tested Applications: Positive Controls:

IF, IP, WB, ELISA WB: U-937 cells, HeLa cells, HepG2 cells, NIH/3T3

Cited Applications: cells, HL-60 cells, SGC-7901 cells IHC, IP, WB

Species Specificity:
human, mouse

IP: U-937 cells,
IF: HeLa cells,

Cited Species: human

## **Background Information**

The galectins, formerly known as S-type lectins, are a family of  $\beta$ -galactoside-binding proteins implicated in modulating cell-cell and cell-matrix interactions. Galectin-9 (LGALS9) is a tandem repeat-type member of the galectin family. It has three isoforms (named galectin-9L, galectin-9M, and galectin-9S): long type of 355 amino acids, medium type of 323 amino acids, and short type of 311 amino acids. Galectin-9 is ubiquitously expressed in a variety of tissues, including lymph nodes and spleen, and overexpressed in Hodgkin disease tissue. It is involved in chemoattraction, apoptosis, and the regulation of cell differentiation and has anti-allergic effects. It has been reported that galectin-9 is a ligand for Tim-3, through which galectin-9 can induce T-helper type 1 lymphocyte (Th1) death.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Junyu Fan	30223894	Stem Cell Res Ther	WB
Jiaxing Yang	24919464	Int J Oncol	
Qi Zhang	36435292	J Nutr Biochem	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




WB result of Galectin 9 antibody (17938-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Galectin 9 transfected NIH/3T3 cells.



IP result of anti-Galectin-9(IP:17938-1-AP, 4ug; Detection:17938-1-AP 1:1500) with U-937 cells lysate 2240 ug.



U-937 cells were subjected to SDS PAGE followed by western blot with 17938-1-AP (Galectin 9 antibody at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 17938-1-AP (Galectin 9 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).