

FURIN Polyclonal antibody

Catalog Number: 18413-1-AP

Featured Product

18 Publications

Basic Information

Catalog Number:

18413-1-AP

Size:

150ul, Concentration: 600 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG13261

GenBank Accession Number:

BC012181

GeneID (NCBI):

5045

Full Name:

furin (paired basic amino acid cleaving enzyme)

Calculated MW:

87 kDa

Observed MW:

90-110 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 µg for 1.0-3.0 mg of total

protein lysate

IHC 1:20-1:200

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human

Cited Species:

human, chicken, rat, mouse

Positive Controls:

WB: HepG2 cells, COLO 320 cells, HeLa cells

IP: HeLa cells,

IHC: human placenta tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

FURIN, also named FUR, PACE, PCSK3, SPC1, and Kex2p, is likely to represent the ubiquitous endoprotease activity within constitutive secretory pathways and capable of cleavage at the RX(K/R)R consensus motif. Furin is synthesized as an inactive zymogen that may minimize the occurrence of premature enzymatic activity that would lead to alternative protein activation or degradation. Glycosylation and other post-translational modifications can make the pre-pro furin run at 100-110 kDa, the mature furin at 98-95 kDa and the shed furin at 90 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Carme Aman	35619054	BMC Genomics	WB
Charles N S Allen	35418836	Front Cell Neurosci	WB
Youli Zhang	28259973	Int J Oncol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of FURIN antibody (18413-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FURIN transfected HepG2 cells.



IP result of anti-FURIN(IP:18413-1-AP, 4ug; Detection:18413-1-AP 1:4000) with HeLa cells lysate 2000 ug.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 18413-1-AP (FURIN Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 18413-1-AP (FURIN Antibody) at dilution of 1:200 (under 40x lens).