For Research Use Only

## CD36 Polyclonal antibody Catalog Number: 18836-1-AP Featured Product

Featured Product

205 Publications




Basic Information	Catalog Number: 18836-1-AP	GenBank Accession Number: BC 008406	Purification Metho Antigen affinity p		
	Size: 150ul , Concentration: 900 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI):	Recommended Dilutions:		
		948	WB 1:500-1:2000 IHC 1:400-1:1000		
		UNIPROT ID: P16671	THC 1:400-1:1000		
		Full Name: CD36 molecule (thrombospondin receptor)			
		Calculated MW: 472 aa, 53 kDa			
		Observed MW: 88 kDa			
Applications	Tested Applications: WB, IHC, ELISA		Positive Controls:		
	Cited Applications:		n peripheral blood pla		
	WB, IHC, IP, CoIP	ITL: numan spleen tissue, num		n liver tissue, humar	
	Species Specificity: human				
	Cited Species: human, pig, bovine, goat Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	retrieval may be performed w	<b>With citrate</b> is an 88-kDa membrane glycoprotei reral tumor cell lines. CD36 binds to and may function in the transport a n as a cell adhesion molecule. It med Mutation of CD36 will cause platele	collagen, thrombospo nd/or as a regulator of diates cytoadherence o et glycoprotein IV defic	ndin, anionic fatty acid transport of Plasmodium ciency which known	
	retrieval may be performed w buffer pH 6.0 CD36, also named as GP3B and GP4, precursors, endothelial cells, and sev phospholipids, long chain fatty acids and oxidized LDL CD36 may functior falciparum parasitized erythrocytes. as CD36 deficiency. Genetic variatio (CHDS7).	is an 88-kDa membrane glycoprotei reral tumor cell lines. CD36 binds to and may function in the transport a n as a cell adhesion molecule. It med Mutation of CD36 will cause platele ns in CD36 are associated with susce	collagen, thrombospo nd/or as a regulator of diates cytoadherence o et glycoprotein IV defic	ndin, anionic fatty acid transport of Plasmodium ciency which known leart disease type 7	
	retrieval may be performed w buffer pH 6.0 CD36, also named as GP3B and GP4, precursors, endothelial cells, and sev phospholipids, long chain fatty acids and oxidized LDL CD36 may function falciparum parasitized erythrocytes. as CD36 deficiency. Genetic variation (CHDS7). Author Pub	is an 88-kDa membrane glycoprotei eral tumor cell lines. CD36 binds to and may function in the transport a nas a cell adhesion molecule. It mer Mutation of CD36 will cause platele ns in CD36 are associated with susce med ID Journal	collagen, thrombospo nd/or as a regulator of diates cytoadherence o et glycoprotein IV defic	ndin, anionic fatty acid transport of Plasmodium ciency which known neart disease type 7 Application	
Background Information	retrieval may be performed w buffer pH 6.0 CD36, also named as GP3B and GP4, precursors, endothelial cells, and sev phospholipids, long chain fatty acids and oxidized LDL CD36 may functior falciparum parasitized erythrocytes. as CD36 deficiency. Genetic variatio (CHDS7). Author Pub Linfeng Ma 330	is an 88-kDa membrane glycoprotei eral tumor cell lines. CD36 binds to and may function in the transport a nas a cell adhesion molecule. It med Mutation of CD36 will cause platele ns in CD36 are associated with susce med ID Journal 07364 Toxicology	collagen, thrombospo nd/or as a regulator of diates cytoadherence o et glycoprotein IV defic	ndin, anionic fatty acid transport of Plasmodium ciency which known neart disease type 7 Application WB	
	retrieval may be performed w buffer pH 6.0CD36, also named as GP3B and GP4, precursors, endothelial cells, and sev phospholipids, long chain fatty acids and oxidized LDL CD36 may function falciparum parasitized erythrocytes. as CD36 deficiency. Genetic variation (CHDS7).AuthorPub Linfeng Ma3300 Meng Zhang289	is an 88-kDa membrane glycoprotei reral tumor cell lines. CD36 binds to and may function in the transport a nas a cell adhesion molecule. It mer Mutation of CD36 will cause platele ns in CD36 are associated with susce med ID Journal 07364 Toxicology 61281 PLoS One	collagen, thrombospo nd/or as a regulator of diates cytoadherence o et glycoprotein IV defic	ndin, anionic fatty acid transport of Plasmodium ciency which known neart disease type 7 Application	
	retrieval may be performed w buffer pH 6.0CD36, also named as GP3B and GP4, precursors, endothelial cells, and sev phospholipids, long chain fatty acids and oxidized LDL CD36 may function falciparum parasitized erythrocytes. as CD36 deficiency. Genetic variation (CHDS7).AuthorPubLinfeng Ma3300Meng Zhang289	is an 88-kDa membrane glycoprotei reral tumor cell lines. CD36 binds to and may function in the transport a nas a cell adhesion molecule. It mer Mutation of CD36 will cause platele ns in CD36 are associated with susce med ID Journal 07364 Toxicology 61281 PLoS One	collagen, thrombospo nd/or as a regulator of diates cytoadherence et glycoprotein IV defic eptibility to coronary h	ndin, anionic fatty acid transport of Plasmodium ciency which known leart disease type 7 Application WB WB,IHC	
Notable Publications	retrieval may be performed w buffer pH 6.0CD36, also named as GP3B and GP4, precursors, endothelial cells, and sev phospholipids, long chain fatty acids and oxidized LDL CD36 may function falciparum parasitized erythrocytes. as CD36 deficiency. Genetic variation (CHDS7).AuthorPubLinfeng Ma3300Meng Zhang289	is an 88-kDa membrane glycoprotei eral tumor cell lines. CD36 binds to and may function in the transport a has a cell adhesion molecule. It mee Mutation of CD36 will cause platele ns in CD36 are associated with susce med ID Journal 07364 Toxicology 61281 PLoS One 03464 Evid Based Comple	collagen, thrombospo nd/or as a regulator of diates cytoadherence et glycoprotein IV defic eptibility to coronary h	ndin, anionic fatty acid transport of Plasmodium ciency which known leart disease type 7 Application WB WB,IHC	
	retrieval may be performed w buffer pH 6.0 CD36, also named as GP3B and GP4, precursors, endothelial cells, and sev phospholipids, long chain fatty acids and oxidized LDL CD36 may functior falciparum parasitized erythrocytes. as CD36 deficiency. Genetic variatio (CHDS7). Author Pub Linfeng Ma 330 Meng Zhang 289 Shuyun Yang 346 Storage: Storage Star 20°C. Stable for one year after Storage Buffer:	is an 88-kDa membrane glycoprotei eral tumor cell lines. CD36 binds to and may function in the transport a has a cell adhesion molecule. It med Mutation of CD36 will cause platele ns in CD36 are associated with susce med ID Journal 07364 Toxicology 61281 PLoS One 03464 Evid Based Comple er shipment.	collagen, thrombospo nd/or as a regulator of diates cytoadherence et glycoprotein IV defic eptibility to coronary h	ndin, anionic fatty acid transport of Plasmodium ciency which known leart disease type 7 Application WB WB,IHC	

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com


Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 18836-1-AP (CD36 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 18836-1-AP (CD36 antibody) at dilution of 1:800 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 18836-1-AP (CD36 antibody) at dilution of 1:800 (under 40x lens).