For Research Use Only

## HMCN1 Polyclonal antibody Catalog Number: 18837-1-AP 4 Publications


oroteintech Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 18837-1-AP	GenBank Accession Number: NM_031935	Purification Method: Antigen affinity purification	
	Size: 150ul, Concentration: 700 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GeneID (NCBI): 83872 UNIPROT ID: Q96RW7	Recommended Dilutions: WB 1:500-1:2000 IHC 1:20-1:200 IF/ICC 1:50-1:500	
		Full Name: hemicentin 1		
		Calculated MW: 613 kDa, 600 kDa, 60 kDa		
		Observed MW: 60 kDa		
Applications	Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls:		
	Cited Applications:	WB : mouse skin tissue, IHC : human colon tissue,		
	WB, IHC, IF			
	Species Specificity: human, mouse, rat	IF/ICC : hTERT-RPE1 cells, MCF-7 cells		
	Cited Species: human, mouse, rat			
	Note-IHC: suggested antigen ı TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen		
Background Information	modules, and a fibulin carboxy-term with age-related macular degenerati	n Willebrand A domain, tandem ir inal module (PMID: 24912920). Mu on. Three alternative transcripts e nst a synthesized peptide correspo	lular protein of the immunoglobulin nmunoglobulin modules, tandem EGF Itations of HMCN1 gene may be associated encoding different HMCN1 isoforms have bee anding to 960-976 amino acids of human	
	superfamily. HMCN1 consists of a vo modules, and a fibulin carboxy-term with age-related macular degenerati described. This antibody raised again HMCN1 can recognize all three isofor	n Willebrand A domain, tandem ir inal module (PMID: 24912920). Mu on. Three alternative transcripts e nst a synthesized peptide correspo	nmunoglobulin modules, tandem EGF utations of HMCN1 gene may be associated ncoding different HMCN1 isoforms have bee	
	superfamily. HMCN1 consists of a vo modules, and a fibulin carboxy-term with age-related macular degenerati described. This antibody raised again HMCN1 can recognize all three isofor Author Put	n Willebrand A domain, tandem ir inal module (PMID: 24912920). Mu on. Three alternative transcripts e ist a synthesized peptide correspo ms of HMCN1.	nmunoglobulin modules, tandem EGF utations of HMCN1 gene may be associated incoding different HMCN1 isoforms have bee inding to 960-976 amino acids of human Application	
Background Information	superfamily. HMCN1 consists of a vo modules, and a fibulin carboxy-term with age-related macular degenerati described. This antibody raised again HMCN1 can recognize all three isofor Author Put Junyu Han 360	n Willebrand A domain, tandem ir inal module (PMID: 24912920). Mu on. Three alternative transcripts e ist a synthesized peptide correspo ms of HMCN1.	nmunoglobulin modules, tandem EGF utations of HMCN1 gene may be associated incoding different HMCN1 isoforms have bee inding to 960-976 amino acids of human Application	
	superfamily. HMCN1 consists of a vo modules, and a fibulin carboxy-term with age-related macular degenerati described. This antibody raised again HMCN1 can recognize all three isofor Author Put Junyu Han 360 Shir Bergson 399	n Willebrand A domain, tandem ir inal module (PMID: 24912920). Mu on. Three alternative transcripts e ist a synthesized peptide correspo ms of HMCN1. med ID Journal 034861 Front Pharmacol	nmunoglobulin modules, tandem EGF utations of HMCN1 gene may be associated incoding different HMCN1 isoforms have bee inding to 960-976 amino acids of human Application WB	
	superfamily. HMCN1 consists of a vo modules, and a fibulin carboxy-term with age-related macular degenerati described. This antibody raised again HMCN1 can recognize all three isofor Author Put Junyu Han 360 Shir Bergson 399	n Willebrand A domain, tandem ir inal module (PMID: 24912920). Mu on. Three alternative transcripts enst a synthesized peptide correspondent rss of HMCN1. med ID Journal 176600 J Exp Med 176600 J Exp Med 176600 PLoS Pathog er shipment. % glycerol, pH7.3	nmunoglobulin modules, tandem EGF stations of HMCN1 gene may be associated incoding different HMCN1 isoforms have been inding to 960-976 amino acids of human Application WB IF	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon using 18837-1-AP (HMCN1 antibody) at dilution of 1:50 (under 10x lens).



 $150kd \rightarrow$   $100kd \rightarrow$   $70kd \rightarrow$   $50kd \rightarrow$   $40kd \rightarrow$  $30kd \rightarrow$ 

Immunohistochemical analysis of paraffinembedded human colon using 18837-1-AP (HMCN1 antibody) at dilution of 1:50 (under 40x lens). mouse skin tissue were subjected to SDS PAGE followed by western blot with 18837-1-AP (HMCN1 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed hTERT-RPE1 cells using HMCN1 antibody (18837-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).