For Research Use Only

MYOD1 Polyclonal antibody Catalog Number: 18943-1-AP Featured Product 11

Featured Product 117 Publications




	Catalog Number: 18943-1-AP	GenBank Accession M BC064493	lumber:	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):		Recommended Dilutions:	
	150ul , Concentration: 600 ug/ml by Nanodrop;	4654 UNIPROT ID: P15172		WB 1:1000-1:6000 IHC 1:50-1:500 IF/ICC 1:400-1:1600	
					Source:
	Rabbit				Full Name: myogenic differentiation 1 Calculated MW: 320 aa, 35 kDa
	Isotype:				
	IgG Immunogen Catalog Number: AG13512				
		Observed MW: 35-45 kDa			
	Applications	WB, IHC, IF/ICC, ELISA WB : mous Cited Applications: skeletal m		Positive Cont	rols:
				eart tissue, human heart tissue, mouse :le tissue, rat heart tissue	
WB, IHC, IF, ChIP			IHC : mouse e	mouse embryo tissue,	
Species Specificity: IF/ICC human, mouse, rat		IF/ICC : HeLa	cells,		
Cited Species: human, mouse, rat, pig, rabbit, chicken, zebrafish, sheep, goat, duck					
Note-IHC: suggested antigen (TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0		vely, antigen			
	MYOD1, also named as BHLHC1 or MYF3, is a 320 amino acid protein, which promotes the transcriptional activity of MYOD1 through its CDK9-mediated phosphorylation. This phosphorylation promotes its function in muscle differentiation. MYOD1 acts as a transcriptional activator that promotes transcription of muscle-specific target gene and plays a role in muscle differentiation. MYOD1 together with MYF5 and MYOG, co-occupies muscle-specific gene promoter core region during myogenesis. Induces fibroblasts to differentiate into myoblasts. he calcualted molecular weight of MYOD1 is 34 kDa, but modified MYOD1 is about 45 kDa. (PMID: 12037670)				
Background Information	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen	scriptional activator thation. MYOD1 together esis. Induces fibroblas	with MYF5 and ts to differentiat	nscription of muscle-specific target gen MYOG, co-occupies muscle-specific gen e into myoblasts. he calcualted	
	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen molecular weight of MYOD1 is 34 kD	scriptional activator thation. MYOD1 together esis. Induces fibroblas	with MYF5 and ts to differentiat 1 is about 45 kD	nscription of muscle-specific target gen MYOG, co-occupies muscle-specific gen e into myoblasts. he calcualted	
	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen molecular weight of MYOD1 is 34 kD Author Pu	scriptional activator th ation. MYOD1 together esis. Induces fibroblas a, but modified MYOD bmed ID Jour	with MYF5 and ts to differentiat 1 is about 45 kD	nscription of muscle-specific target gen MYOG, co-occupies muscle-specific gen e into myoblasts. he calcualted a. (PMID: 12037670)	
	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen molecular weight of MYOD1 is 34 kD Author Pu Yuanyuan Wu 36	scriptional activator th ation. MYOD1 together esis. Induces fibroblas a, but modified MYOD bmed ID Jour 075558 Cell	with MYF5 and ts to differentiat 1 is about 45 kD	nscription of muscle-specific target gen MYOG, co-occupies muscle-specific gen e into myoblasts. he calcualted a. (PMID: 12037670) Application	
	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen molecular weight of MYOD1 is 34 kD Author Pu Yuanyuan Wu 36 Peng Ren 36	scriptional activator th ation. MYOD1 together esis. Induces fibroblas a, but modified MYOD bmed ID Jour 075558 Cell 118887 Fror	with MYF5 and ts to differentiat 1 is about 45 kD mal Signal	nscription of muscle-specific target gen MYOG, co-occupies muscle-specific gen e into myoblasts. he calcualted a. (PMID: 12037670) Application WB	
	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen molecular weight of MYOD1 is 34 kD Author Pu Yuanyuan Wu 36 Peng Ren 36	scriptional activator th ation. MYOD1 together esis. Induces fibroblas a, but modified MYOD bmed ID Jour 075558 Cell 118887 Fror	with MYF5 and ts to differentian 1 is about 45 kD nal Signal tt Genet	nscription of muscle-specific target gen MYOG, co-occupies muscle-specific gen e into myoblasts. he calcualted a. (PMID: 12037670) Application WB IHC	
Notable Publications	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen molecular weight of MYOD1 is 34 kD Author Pu Yuanyuan Wu 36 Peng Ren 36 Hongyi Zhou 34 Storage: Store at -20°C. Stable for one year aff Storage Buffer:	scriptional activator thation. MYOD1 together esis. Induces fibroblas a, but modified MYOD bmed ID Jour 075558 Cell 118887 Fror 502418 Int J	with MYF5 and ts to differentian 1 is about 45 kD nal Signal tt Genet	nscription of muscle-specific target gen MYOG, co-occupies muscle-specific gen e into myoblasts. he calcualted a. (PMID: 12037670) Application WB IHC	
Notable Publications	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen molecular weight of MYOD1 is 34 kD Author Pu Yuanyuan Wu 36 Peng Ren 36 Hongyi Zhou 34 Storage: Store at -20°C. Stable for one year after	scriptional activator thation. MYOD1 together esis. Induces fibroblas a, but modified MYOD bmed ID Jour 075558 Cell 118887 Fror 502418 Int J ser shipment.	with MYF5 and ts to differentian 1 is about 45 kD nal Signal tt Genet	nscription of muscle-specific target gen MYOG, co-occupies muscle-specific gen e into myoblasts. he calcualted a. (PMID: 12037670) Application WB IHC	
Background Information Notable Publications Storage *** 20ul sizes contain 0.1% BSA	differentiation. MYOD1 acts as a tran and plays a role in muscle differentia promoter core region during myogen molecular weight of MYOD1 is 34 kD Author Pu Yuanyuan Wu 366 Peng Ren 366 Hongyi Zhou 344 Storage: Stora at -20°C. Stable for one year affection Storage Buffer: PBS with 0.02% sodium azide and 500	scriptional activator thation. MYOD1 together esis. Induces fibroblas a, but modified MYOD bmed ID Jour 075558 Cell 118887 Fror 502418 Int J ser shipment.	with MYF5 and ts to differentian 1 is about 45 kD nal Signal tt Genet	Application MYOG, co-occupies muscle-specific target gen into myoblasts. he calcualted (PMID: 12037670) Application WB IHC	

chnical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





mouse heart tissue were subjected to SDS PAGE followed by western blot with 18943-1-AP (MYOD1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded mouse embryo tissue slide using 18943-1-AP (MYOD1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MYOD1 antibody (18943-1-AP) at dilution of 1:800 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red).