For Research Use Only

CD4 Polyclonal antibody Catalog Number: 19068-1-AP 33 Publications




Basic Information	Catalog Number: 19068-1-AP	GenBank Accession Number: BC025782	Purification Method: Antigen affinity purification	
	Size: 150ul , Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG13616	GenelD (NCBI): 920	Recommended Dilutions: WB 1:500-1:3000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
				ENSEMBL Gene ID: ENSG00000010610
		UNIPROT ID:		
		P01730		
		Full Name: CD4 molecule		
		Calculated MW:		
		55 kDa		
		Observed MW: 55 kDa		
		Applications	Tested Applications:	Positive C
WB, IP, ELISA Cited Applications:				
WB	tissue			
Species Specificity:	IP : mouse thymus tissue,			
	Species Specificity.			
	human, mouse, rat, pig			
	human, mouse, rat, pig Cited Species:			
Background Informatior	human, mouse, rat, pig Cited Species: human, mouse, rat CD4 is a 55-kDa transmembrane glymacrophages, and dendritic cells. CD	94 is an accessory protein for MHC o	lls, majority of thymocytes, monocytes, class-II antigen/T-cell receptor interaction D4 serves as a receptor for the human	
	human, mouse, rat, pig Cited Species: human, mouse, rat CD4 is a 55-kDa transmembrane gly macrophages, and dendritic cells. CD plays an important role in T helper co immunodeficiency virus (HIV).	04 is an accessory protein for MHC c ell development and activation. Cl	lass-II antigen/T-cell receptor interaction D4 serves as a receptor for the human	
	human, mouse, rat, pig Cited Species: human, mouse, rat CD4 is a 55-kDa transmembrane gly macrophages, and dendritic cells. CD plays an important role in T helper co immunodeficiency virus (HIV). Author Put	04 is an accessory protein for MHC of ell development and activation. Cl omed ID Journal	class-II antigen/T-cell receptor interaction	
	human, mouse, rat, pig Cited Species: human, mouse, rat CD4 is a 55-kDa transmembrane glyy macrophages, and dendritic cells. CD plays an important role in T helper of immunodeficiency virus (HIV). Author Put Qicai Liu 315	04 is an accessory protein for MHC c ell development and activation. CI pmed ID Journal 521106 Mol Med	lass-II antigen/T-cell receptor interaction D4 serves as a receptor for the human	
Background Informatior Notable Publications	human, mouse, rat, pig Cited Species: human, mouse, rat CD4 is a 55-kDa transmembrane gly macrophages, and dendritic cells. CD plays an important role in T helper co immunodeficiency virus (HIV). Author Put Qicai Liu 311 Tsai-Wei Hung 342	04 is an accessory protein for MHC of ell development and activation. Cl omed ID Journal	lass-II antigen/T-cell receptor interaction D4 serves as a receptor for the human	
	human, mouse, rat, pig Cited Species: human, mouse, rat CD4 is a 55-kDa transmembrane gly macrophages, and dendritic cells. CD plays an important role in T helper co immunodeficiency virus (HIV). Author Put Qicai Liu 311 Tsai-Wei Hung 342	04 is an accessory protein for MHC c ell development and activation. Cl omed ID Journal 521106 Mol Med 576050 Int J Mol Sci	Lass-II antigen/T-cell receptor interaction D4 serves as a receptor for the human Application	
	human, mouse, rat, pig Cited Species: human, mouse, rat CD4 is a 55-kDa transmembrane gly macrophages, and dendritic cells. CD plays an important role in T helper co immunodeficiency virus (HIV). Author Put Qicai Liu 311 Tsai-Wei Hung 342	04 is an accessory protein for MHC of ell development and activation. Cl omed ID Journal 521106 Mol Med 576050 Int J Mol Sci 505457 Front Immunol ter shipment.	Lass-II antigen/T-cell receptor interaction D4 serves as a receptor for the human Application	
Notable Publications	human, mouse, rat, pig Cited Species: human, mouse, rat CD4 is a 55-kDa transmembrane glyy macrophages, and dendritic cells. CD plays an important role in T helper co immunodeficiency virus (HIV). Author Put Qicai Liu 31! Tsai-Wei Hung 34! Yulin Deng 36! Storage: Storage at -20°C. Stable for one year affectors	04 is an accessory protein for MHC of ell development and activation. Cl 5000000000000000000000000000000000000	Lass-II antigen/T-cell receptor interaction D4 serves as a receptor for the human Application	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



U-937 cells were subjected to SDS PAGE followed by western blot with 19068-1-AP (CD4 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.

 $250 \text{ kDa} \rightarrow$ $150 \text{ kDa} \rightarrow$

 $100 \text{ kDa} \rightarrow$

70 kDa \rightarrow 50 kDa \rightarrow 40 kDa \rightarrow

30 kDa-

20 kDa→ 15 kDa→

> IP result of anti-CD4 (IP:19068-1-AP, 4ug; Detection:19068-1-AP 1:500) with mouse thymus tissue lysate 2800ug.