For Research Use Only

ATG14/Barkor (N-terminal) Polyclonal antibody

www.ptglab.com

Catalog Number: 19491-1-AP

Featured Product

36 Publications

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

Antigen affinity purification

19491-1-AP Size:

NM_014924 GeneID (NCBI):

Recommended Dilutions:

150ul , Concentration: 350 ug/ml by

Rabbit

Isotype:

IgG

22863

WB: 1:500-1:1000 IHC: 1:50-1:500 IF/ICC: 1:50-1:500

Nanodrop; Source:

UNIPROT ID: Q6ZNE5

Full Name:

KIAA0831

Calculated MW: 55 kDa

65-68 kDa

Observed MW:

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IF, IP

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat, pig, canine, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, A431 cells, mouse brain tissue,

NIH/3T3 cells

IHC: mouse brain tissue, IF/ICC: MCF-7 cells,

Background Information

Barkor, also named as KIAA0831, Atg14 and Atg14L, is required for both basal and inducible autophagy. It plays a role in autophagosome formation and MAP1LC3/LC3 conjugation to phosphatidylethanolamine. Barkor promotes BECN1 translocation from the trans-Golgi network to autophagosomes. This antibody is specific to KIAA0831(Barkor).

Notable Publications

Author	Pubmed ID	Journal	Application
Luis Muniz-Feliciano	28933590	Autophagy	WB
Qiang Fu	25234643	J Gen Virol	WB
Xueqiang Peng	32949647	Biochim Biophys Acta Mol Cell Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:50 and CoraLite488-Conjugated Goat Anti-Rabbit IgG(H+L).