For Research Use Only

ND2 Polyclonal antibody Catalog Number: 19704-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: 19704-1-AP	GenBank Accession Number: GQ 398480	Purification Method: Antigen affinity purification
	Size: 150ul, Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI):	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500
		4536	
		UNIPROT ID:	
		P03891	
		Full Name: NADH dehydrogenase, subunit 2 (complex I)	
		Observed MW:	
		42-44 kDa	
		Applications	Tested Applications:
WB, IHC, ELISA			
Cited Applications: IHC : human sto WB, IHC tissue			nan stomach tissue, human stomach cance
Species Specificity: human, mouse Cited Species: human, mouse, rat			
	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternati retrieval may be performed v buffer pH 6.0	ively, antigen	
Background Information	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternative retrieval may be performed v buffer pH 6.0 ND2, also named as MTND2, NADH2 mitochondrial membrane respirator	ively, antigen vith citrate and ND2, belongs to the complex I y chain NADH dehydrogenase (Cor lysis. Complex I functions in the tra	nplex I) that is believed to belong to the ansfer of electrons from NADH to the
	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternative retrieval may be performed v buffer pH 6.0 ND2, also named as MTND2, NADH2 mitochondrial membrane respirator minimal assembly required for cata respiratory chain. The immediate el	ively, antigen vith citrate and ND2, belongs to the complex I y chain NADH dehydrogenase (Cor lysis. Complex I functions in the tra	ansfer of electrons from NADH to the
Background Information	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternative retrieval may be performed ve buffer pH 6.0 ND2, also named as MTND2, NADH2 mitochondrial membrane respirator minimal assembly required for cata respiratory chain. The immediate el	ively, antigen vith citrate and ND2, belongs to the complex I y chain NADH dehydrogenase (Cor lysis. Complex I functions in the tra ectron acceptor for the enzyme is b	nplex I) that is believed to belong to the ansfer of electrons from NADH to the selieved to be ubiquinone.
	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternati retrieval may be performed v buffer pH 6.0ND2, also named as MTND2, NADH2 mitochondrial membrane respirator minimal assembly required for cata respiratory chain. The immediate elAuthorPu Ying ShuYing Shu36	ively, antigen with citrate and ND2, belongs to the complex I y chain NADH dehydrogenase (Cor lysis. Complex I functions in the tr ectron acceptor for the enzyme is b bmed ID Journal	nplex I) that is believed to belong to the ansfer of electrons from NADH to the believed to be ubiquinone. Application WB
	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternati retrieval may be performed v buffer pH 6.0ND2, also named as MTND2, NADH2 mitochondrial membrane respirator minimal assembly required for cata respiratory chain. The immediate elAuthorPu Ying ShuAlex Gallinat36	ively, antigen vith citrate and ND2, belongs to the complex I y chain NADH dehydrogenase (Cor lysis. Complex I functions in the tra ectron acceptor for the enzyme is b bmed ID Journal 314841 EMBO J	nplex I) that is believed to belong to the ansfer of electrons from NADH to the believed to be ubiquinone. Application WB
	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternati retrieval may be performed v buffer pH 6.0ND2, also named as MTND2, NADH2 mitochondrial membrane respirator minimal assembly required for cata respiratory chain. The immediate elAuthorPu Ying ShuYing Shu36Alex Gallinat36	ively, antigen vith citrate and ND2, belongs to the complex I y chain NADH dehydrogenase (Cor lysis. Complex I functions in the tra ectron acceptor for the enzyme is b bmed ID Journal 314841 EMBO J 341940 Free Radic Biol M 492917 EMBO J	nplex I) that is believed to belong to the ansfer of electrons from NADH to the selieved to be ubiquinone. Application WB Med WB


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



mouse liver tissue were subjected to SDS PAGE followed by western blot with 19704-1-AP (ND2 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 19704-1-AP (ND2 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 19704-1-AP (ND2 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).