For Research Use Only

RIM2-Specific Polyclonal antibody

Catalog Number: 20093-1-AP



Purification Method:

WB 1:500-1:2000

IHC 1:50-1:500

IHC: mouse brain tissue,

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 20093-1-AP NM 001100117

re: GenelD (NCBI):

150ul , Concentration: 500 µg/ml by 9699

Nanodrop and 280 µg/ml by Bradford method using BSA as the standard; regulating sy

method using BSA as the standard; regulating synaptic membrane

Source:exocytosis 2RabbitCalculated MW:Isotype:160 kDaIgGObserved MW:

133 kDa

Applications

Tested Applications: Positive Controls: IHC, WB, ELISA WB: LO2 cells,

Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

RIMS2, also named as KIAA0751, RAB3IP3 and RIM2, is a rab effector which involved in exocytosis. RIMS2 may act as scaffold protein. This antibody is specific to RIMS2. It recognizes all the isoforms except isoform of RIMS2.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



LO2 cells were subjected to SDS PAGE followed by western blot with 20093-1-AP (RIM2-Specific antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20093-1-AP (RIM2-Specific antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20093-1-AP (RIM2-Specific antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).