For Research Use Only

CD9 Polyclonal antibody

Catalog Number:20597-1-AP

Featured Product





Basic Information	Catalog Number: 20597-1-AP	GenBank Accession Number: BC011988	Purification Method: Antigen affinity purification
	Size: 150ul, Concentration: 600 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number:	GeneID (NCBI): 928 UNIPROT ID: P21926 Full Name: CD9 molecule Calculated MW: 228 aa, 25 kDa	Recommended Dilutions: WB: 1:2000-1:12000 IHC: 1:1000-1:4000 IF/ICC: 1:50-1:500
	AG14546	Observed MW: 23-30 kDa	
Applications	Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls: WB : HCT 116 cells, HeLa cells, L02 cells	
	Cited Applications: IHC : human lung cancer tissue, human breast can tissue, human endometrial cancer tissue, human species Specificity: Species Specificity: cancer tissue human IF/ICC : MCF-7 cells, Cited Species: IF/ICC : MCF-7 cells, human, mouse, rat, pig, rabbit, monkey, chicken, zebrafish, goat, bat IF/ICC : MCF-7 cells,		an lung cancer tissue, human breast cancer man endometrial cancer tissue, human ovar
			ICF-7 cells,
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
		ith citrate	
Background Information	buffer pH 6.0 The cell-surface molecule CD9, a me and other membrane proteins, and is enhances membrane fusion betweer often used as a mesenchymal stem of	mber of the transmembrane-4 sup postulated to participate in cell m muscle cells and promotes viral in ell marker (PMID:18005405). The C ains and one N-glycosylation site	D9 antigen appears to be a 227-amino acid (PMID: 1840589). This antibody detects band
	buffer pH 6.0 The cell-surface molecule CD9, a me and other membrane proteins, and is enhances membrane fusion betweer often used as a mesenchymal stem of molecule with four hydrophobic dom of 23-30 kDa, it may be due to the dif	mber of the transmembrane-4 sup postulated to participate in cell m muscle cells and promotes viral in ell marker (PMID:18005405). The C ains and one N-glycosylation site	igration and adhesion. Expression of CD9 nfection in some cells (PMID:10459022). It is CD9 antigen appears to be a 227-amino acid (PMID: 1840589). This antibody detects banc
	buffer pH 6.0 The cell-surface molecule CD9, a me and other membrane proteins, and is enhances membrane fusion between often used as a mesenchymal stem of molecule with four hydrophobic dom of 23-30 kDa, it may be due to the dis	mber of the transmembrane-4 sup postulated to participate in cell m muscle cells and promotes viral in ell marker (PMID:18005405). The C ains and one N-glycosylation site ference of glycosylations (PMID: 8	igration and adhesion. Expression of CD9 nfection in some cells (PMID:10459022). It is D9 antigen appears to be a 227-amino acid (PMID: 1840589). This antibody detects banc 701996). Application
Background Information Notable Publications	buffer pH 6.0 The cell-surface molecule CD9, a meand other membrane proteins, and is enhances membrane fusion betweer often used as a mesenchymal stem comolecule with four hydrophobic dom of 23-30 kDa, it may be due to the difference of the difference o	mber of the transmembrane-4 sup postulated to participate in cell m muscle cells and promotes viral in ell marker (PMID:18005405). The C ains and one N-glycosylation site freence of glycosylations (PMID: 8 med ID Journal	igration and adhesion. Expression of CD9 nfection in some cells (PMID:10459022). It is D9 antigen appears to be a 227-amino acid (PMID: 1840589). This antibody detects banc 701996). Application
	buffer pH 6.0 The cell-surface molecule CD9, a meand other membrane proteins, and is enhances membrane fusion betweer often used as a mesenchymal stem of 23-30 kDa, it may be due to the dimonstration of 23-30 kDa, it may be due to the dimonstrational stem of 23-30 kDa, it may be due to the dimonstratin stem of 23-30 kDa, it may be due to the dimonstration	mber of the transmembrane-4 suppostulated to participate in cell muscle cells and promotes viral in ell marker (PMID:18005405). The Cains and one N-glycosylation site ference of glycosylations (PMID: 8 pmed ID Journal Provide Front Bioeng Biot	igration and adhesion. Expression of CD9 nfection in some cells (PMID:10459022). It is CD9 antigen appears to be a 227-amino acid (PMID: 1840589). This antibody detects band 701996). Application wB
	buffer pH 6.0 The cell-surface molecule CD9, a meand other membrane proteins, and is enhances membrane fusion betweer often used as a mesenchymal stem of 23-30 kDa, it may be due to the dimonstration of 23-30 kDa, it may be due to the dimonstrational stem of 23-30 kDa, it may be due to the dimonstratin stem of 23-30 kDa, it may be due to the dimonstration	mber of the transmembrane-4 suppostulated to participate in cell muscle cells and promotes viral in ell marker (PMID:18005405). The Cains and one N-glycosylation site iference of glycosylations (PMID: 8 med ID Journal 246376 Front Bioeng Biot 596392 Viruses 271212 Nat Commun	ingration and adhesion. Expression of CD9 Infection in some cells (PMID:10459022). It is D9 antigen appears to be a 227-amino acid (PMID: 1840589). This antibody detects band 701996). Application wB WB


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20597-1-AP (CD9 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 20597-1-AP (CD9 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 20597-1-AP (CD9 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using CD9 antibody (20597-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).