For Research Use Only

SOST/Sclerostin Polyclonal antibody

Catalog Number:21933-1-AP

Featured Product

15 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com


Basic Information	Catalog Number: 21933-1-AP	GenBank Accession Number: BC 101086	Purification Method:	
			Antigen affinity purification	
	Size: 150ul , Concentration: 550 ug/ml by	GenelD (NCBI): 50964	Recommended Dilutions: IHC 1:50-1:500	
	Nanodrop;	UNIPROT ID:	IF/ICC 1:50-1:500	
	Source:	Q9BQB4		
	Rabbit	Full Name:		
	lsotype:	sclerosteosis		
	IgG	Calculated MW:		
	Immunogen Catalog Number: AG16567	213 aa, 24 kDa		
Applications	Tested Applications:	Positive Controls:		
	IHC, IF/ICC, ELISA		IHC : human kidney tissue, human breast cancer tissu	
	Cited Applications: WB, IHC, IF	mouse lung tissue, mouse embryo tissue		
	Species Specificity: human, mouse	IF/ICC : HepG2 cells,		
	Cited Species: human, mouse, rat			
	Note-IHC: suggested antigen I TE buffer pH 9.0; (*) Alternati	vely, antigen		
	retrieval may be performed w buffer pH 6.0	vith citrate		
Background Information	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-selec (BMP) antagonists. Loss-of-function n	with a C-terminal cysteine knot-l cted gene aberrative in neuroblas nutations in this gene are associa ive bone overgrowth. A deletion c	toma) family of bone morphogenetic protein ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduce	
	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-selec (BMP) antagonists. Loss-of-function n sclerosteosis, which causes progressi sclerostin expression, is associated w	with a C-terminal cysteine knot-l cted gene aberrative in neuroblas nutations in this gene are associa ive bone overgrowth. A deletion c	ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduced	
	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-selec (BMP) antagonists. Loss-of-function n sclerosteosis, which causes progress sclerostin expression, is associated w	with a C-terminal cysteine knot-l cted gene aberrative in neuroblas nutations in this gene are associa- ive bone overgrowth. A deletion of vith a milder form of the disorder bmed ID Journal	toma) family of bone morphogenetic protein ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduce called van Buchem disease Application	
Background Information Notable Publications	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-selec (BMP) antagonists. Loss-of-function m sclerosteosis, which causes progressis sclerostin expression, is associated w Author Put Fang He	with a C-terminal cysteine knot-l cted gene aberrative in neuroblas nutations in this gene are associa ive bone overgrowth. A deletion c vith a milder form of the disorder	toma) family of bone morphogenetic protein ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduced called van Buchem disease Application	
	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-select (BMP) antagonists. Loss-of-function m sclerosteosis, which causes progressi sclerostin expression, is associated w Author Put Fang He 33 Xiaoan Wei 35	with a C-terminal cysteine knot-l cted gene aberrative in neuroblas nutations in this gene are associa ive bone overgrowth. A deletion of vith a milder form of the disorder bmed ID Journal 159018 Aging (Albany I	toma) family of bone morphogenetic protein ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduce called van Buchem disease Application NY) WB	
	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-select (BMP) antagonists. Loss-of-function m sclerosteosis, which causes progressi sclerostin expression, is associated w Author Put Fang He 33 Xiaoan Wei 35	with a C-terminal cysteine knot-l cted gene aberrative in neuroblas nutations in this gene are associa- ive bone overgrowth. A deletion of vith a milder form of the disorder bmed ID Journal 159018 Aging (Albany I 611810 EMBO Mol Med	toma) family of bone morphogenetic protein ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduce called van Buchem disease Application NY) WB IF	
Notable Publications	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-selec (BMP) antagonists. Loss-of-function m sclerosteosis, which causes progressi sclerostin expression, is associated w Author Put Fang He 33 Xiaoan Wei 35 Marko Lucijanic 29 Storage: Storage Buffer:	with a C-terminal cysteine knot- cted gene aberrative in neuroblas nutations in this gene are associa- ive bone overgrowth. A deletion of vith a milder form of the disorder bmed ID Journal 159018 Aging (Albany I 611810 EMBO Mol Med 532161 Ann Hematol ter shipment.	toma) family of bone morphogenetic protein ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduce called van Buchem disease Application NY) WB IF	
	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-selec (BMP) antagonists. Loss-of-function m sclerosteosis, which causes progressi sclerostin expression, is associated w Author Put Fang He 33 Xiaoan Wei 35 Marko Lucijanic 29 Storage: Storage Buffer: PBS with 0.02% sodium azide and 50	with a C-terminal cysteine knot- cted gene aberrative in neuroblas nutations in this gene are associa- ive bone overgrowth. A deletion of vith a milder form of the disorder bmed ID Journal 159018 Aging (Albany I 611810 EMBO Mol Med 532161 Ann Hematol ter shipment.	toma) family of bone morphogenetic protein ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduce called van Buchem disease Application NY) WB IF	
Notable Publications	buffer pH 6.0 Sclerostin is a secreted glycoprotein the DAN (differential screening-selec (BMP) antagonists. Loss-of-function m sclerosteosis, which causes progressi sclerostin expression, is associated w Author Put Fang He 33 Xiaoan Wei 35 Marko Lucijanic 29 Storage: Storage Buffer:	with a C-terminal cysteine knot- cted gene aberrative in neuroblas nutations in this gene are associa- ive bone overgrowth. A deletion of vith a milder form of the disorder bmed ID Journal 159018 Aging (Albany I 611810 EMBO Mol Med 532161 Ann Hematol ter shipment.	toma) family of bone morphogenetic protein ted with an autosomal-recessive disorder, lownstream of this gene, which causes reduce called van Buchem disease Application NY) WB IF	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffin-21933- embedded human kidney tissue slide using 21933-00 1-AP (Sclerostin antibody) at dilution of 1:200 rieval (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Sclerostin antibody (21933-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).

Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 21933-1-AP (Sclerostin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).