For Research Use Only

CCL17 Polyclonal antibody

Catalog Number:22342-1-AP 2 Publications



Basic Information

Catalog Number: 22342-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

Source:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 300 ug/ml by

BC112066

IHC 1:50-1:500

Nanodrop and 220 ug/ml by Bradford $\ensuremath{\,\,{\sf UNIPROT\,ID:}}$ method using BSA as the standard;

Q92583 Full Name:

Rabbit chemokine (C-C motif) ligand 17

Isotype: Calculated MW: 94 aa, 11 kDa

Immunogen Catalog Number:

AG17832

Applications

Tested Applications:

IHC FIISA

Cited Applications:

IHC

Species Specificity:

human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human colon tissue, human placenta tissue,

human colon cancer tissue

Background Information

CCL17, also named as TARC, is a member of the CC chemokine family, and is highly expressed by thymus and other cells, including keratinocytes, endothelial cells, dendritic cells, bronchial epithelial cells and fibroblasts. CCL17 $plays\ important\ roles\ in\ T\ cell\ development\ in\ thymus\ as\ well\ as\ in\ trafficking\ and\ activation\ of\ mature\ T\ cells.$ CCL17 induced chemotaxis and Ca2+ influx in the cells stably expressing CCR4, thereby demonstrating the ligandreceptor relationship between CCL17 and CCR4.

Notable Publications

Author	Pubmed ID	Journal	Application
Shoko Nakayama	27072191	Br J Haematol	IHC
Emma Kroeze	38948925	Hemasphere	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 22342-1-AP (CCL17 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 22342-1-AP (CCL17 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).