For Research Use Only

## SOX7 Polyclonal antibody Catalog Number:23925-1-AP 5 Publications





Basic Information	Catalog Number: 23925-1-AP	GenBank Accession Number: BC071947	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul , Concentration: 750 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG10323	83595	WB 1:500-1:1000
		UNIPROT ID:	IHC 1:50-1:500 IF/ICC 1:200-1:800
		Q9BT81	111100 1.200 1.000
		Full Name: SRY (sex determining region Y)-box 7 Calculated MW:	
		388 aa, 42 kDa	
		Observed MW:	
		42-45 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, IF/ICC, ELISA	WB:CC	DLO 320 cells, mouse heart tissue, PC-3 cells
	Cited Applications: WB, IF, IP	NB, IF, IP     IFC : numan prostate cancer tissue, numan colon tissu       Species Specificity:     IF/ICC : U2OS cells,	
	Species Specificity: human, mouse		
	Cited Species:		
	human, mouse		
	Note-IHC: suggested antigen retrieval with <b>TE buffer pH 9.0; (*)</b> Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Information	SOX7, also named as Transcription factor SOX-7, is a 388 amino acid protein, which is widely expressed in adult and fetal tissues and tends to be down-regulated in prostate adenocarcinomas and colorectal tumors due to promoter hypermethylation. SOX7 binds to and activates the CDH5 promoter, hence plays a role in the transcriptional regulation of genes expressed in the hemogenic endothelium and blocks further differentiation into blood precursors. SOX7 may be required for the survival of both hematopoietic and endothelial precursors during specification.		
	precursors. SOX7 may be required for	-	
	precursors. SOX7 may be required for specification.	r the survival of both hematopoi	etic and endothelial precursors during
	precursors. SOX7 may be required for specification. Author Pul	r the survival of both hematopoid bmed ID Journal	etic and endothelial precursors during Application
Notable Publications	precursors. SOX7 may be required for specification.           Author         Pul           Baolei Li         352	r the survival of both hematopoid bmed ID Journal 260939 Mol Genet Gen	etic and endothelial precursors during Application nomics WB,IP
	precursors. SOX7 may be required for specification.           Author         Pul           Baolei Li         352	r the survival of both hematopoid bmed ID Journal	etic and endothelial precursors during Application
	precursors. SOX7 may be required for specification. Author Pul Baolei Li 352 Qiao-Yang Sun 312	r the survival of both hematopoid bmed ID Journal 260939 Mol Genet Gen	etic and endothelial precursors during Application wmics WB,IP WB
	precursors. SOX7 may be required for specification. Author Pul Baolei Li 35: Qiao-Yang Sun 31:	r the survival of both hematopoid bmed ID Journal 260939 Mol Genet Gen 332289 Oncogene 846290 Cell Death Dis ter shipment.	etic and endothelial precursors during Application wmics WB,IP WB
Notable Publications	precursors. SOX7 may be required for specification. Author Pul Baolei Li 35: Qiao-Yang Sun 31: Nanchao Hong 33: Storage: Storage Suffer:	r the survival of both hematopoid bmed ID Journal 260939 Mol Genet Gen 332289 Oncogene 846290 Cell Death Dis ter shipment.	etic and endothelial precursors during Application wmics WB,IP WB


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data








Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 23925-1-AP (SOX7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 23925-1-AP (SOX7 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using SOX7 antibody (23925-1-AP) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594phalloidin (red).