For Research Use Only

GFAP Polyclonal antibody Catalog Number:23935-1-AP Featured Product






Basic Information	Catalog Number: 23935-1-AP	GenBank Accession Number: BC013596	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 800 ug/ml by	2670	WB: 1:5000-1:50000	
	Nanodrop;	UNIPROT ID:	IP: 0.5-4.0 ug for 1.0-3.0 mg of total	
	Source:	P14136	protein lysate IHC: 1:200-1:800	
	Rabbit	Full Name:	IF-P: 1:500-1:2000	
	Isotype:	glial fibrillary acidic protein	IF-Fro: 1:50-1:500	
	lgG	Calculated MW:		
	Immunogen Catalog Number: AG20853	432 aa, 50 kDa		
		Observed MW: 45-50 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF-P, IF-Fro, IP, ELISA	WB : U-251 cells, mouse brain tissue, rat brain tissue		
	Cited Applications:	IP : mouse brain tissue,		
	WB, IHC, IF	IHC : m	ouse brain tissue, human gliomas tissue	
	Species Specificity: human, mouse, rat	IF-P : ra	IF-P : rat brain tissue,	
	Cited Species:	IF-Fro:	mouse brain tissue,	
	human, mouse, rat, macaque		······································	
	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen		
	GFAP (Glial fibrillary acidic protein) is a type III intermediate filament (IF) protein specific to the central nervous system (CNS). GFAP is one of the main components of the intermediate filament network in astrocytes and has bee proposed as playing a role in cell migration, cell motility, maintaining mechanical strength, and in mitosis. GFAP is expressed in central nervous system cells, predominantly in astrocytes. GFAP is commonly used as an astrocyte marker. However, GFAP is also present in peripheral glia and in non-CNS cells, including fibroblasts, chondrocytes, lymphocytes, and liver stellate cells (PMID: 21219963).			
Background Information	system (CNS). GFAP is one of the ma proposed as playing a role in cell mi expressed in central nervous system marker. However, GFAP is also prese	in components of the intermedia gration, cell motility, maintaini cells, predominantly in astrocyunt int in peripheral glia and in non-	ate filament network in astrocytes and has bee ng mechanical strength, and in mitosis. GFAP is tes. GFAP is commonly used as an astrocyte	
	system (CNS). GFAP is one of the ma proposed as playing a role in cell mi expressed in central nervous system marker. However, GFAP is also prese lymphocytes, and liver stellate cells	in components of the intermedia gration, cell motility, maintaini cells, predominantly in astrocyu nt in peripheral glia and in non- (PMID: 21219963).	ate filament network in astrocytes and has beer ng mechanical strength, and in mitosis. GFAP is tes. GFAP is commonly used as an astrocyte CNS cells, including fibroblasts, chondrocytes,	
	system (CNS). GFAP is one of the ma proposed as playing a role in cell mi expressed in central nervous system marker. However, GFAP is also prese lymphocytes, and liver stellate cells Author Pu	in components of the intermedia gration, cell motility, maintaini cells, predominantly in astrocy int in peripheral glia and in non- (PMID: 21219963).	ate filament network in astrocytes and has been ng mechanical strength, and in mitosis. GFAP is tes. GFAP is commonly used as an astrocyte CNS cells, including fibroblasts, chondrocytes, Application	
Background Information	system (CNS). GFAP is one of the maproposed as playing a role in cell mi expressed in central nervous system marker. However, GFAP is also prese lymphocytes, and liver stellate cells Author Pu Pengyi Zhou 36	in components of the intermedia gration, cell motility, maintaini cells, predominantly in astrocyu- nt in peripheral glia and in non- (PMID: 21219963).	ate filament network in astrocytes and has beer ng mechanical strength, and in mitosis. GFAP is tes. GFAP is commonly used as an astrocyte CNS cells, including fibroblasts, chondrocytes, Application IF,WB	
	system (CNS). GFAP is one of the maproposed as playing a role in cell mi expressed in central nervous system marker. However, GFAP is also prese lymphocytes, and liver stellate cells Author Pu Pengyi Zhou 36 Shadan S Yarandi 33	in components of the intermedia gration, cell motility, maintaini cells, predominantly in astrocyt int in peripheral glia and in non- (PMID: 21219963). bmed ID Journal 260151 J Mol Histol 2137166 PLoS One	ate filament network in astrocytes and has beer ng mechanical strength, and in mitosis. GFAP is tes. GFAP is commonly used as an astrocyte CNS cells, including fibroblasts, chondrocytes, Application IF,WB IF	
	system (CNS). GFAP is one of the maproposed as playing a role in cell mi expressed in central nervous system marker. However, GFAP is also prese lymphocytes, and liver stellate cells Author Pu Pengyi Zhou 36 Shadan S Yarandi 33	in components of the intermedia gration, cell motility, maintaini cells, predominantly in astrocyu- nt in peripheral glia and in non- (PMID: 21219963).	ate filament network in astrocytes and has been ng mechanical strength, and in mitosis. GFAP is es. GFAP is commonly used as an astrocyte CNS cells, including fibroblasts, chondrocytes, Application IF,WB IF	
Notable Publications	system (CNS). GFAP is one of the maproposed as playing a role in cell mi expressed in central nervous system marker. However, GFAP is also prese lymphocytes, and liver stellate cells Author Pu Pengyi Zhou 36 Shadan S Yarandi 33	in components of the intermedia gration, cell motility, maintaini cells, predominantly in astrocyt int in peripheral glia and in non- (PMID: 21219963). bmed ID Journal 260151 J Mol Histol 2137166 PLoS One 2819195 Aging (Alban) ter shipment.	ate filament network in astrocytes and has been ng mechanical strength, and in mitosis. GFAP is tes. GFAP is commonly used as an astrocyte CNS cells, including fibroblasts, chondrocytes, Application IF,WB IF	
	system (CNS). GFAP is one of the maproposed as playing a role in cell mi expressed in central nervous system marker. However, GFAP is also prese lymphocytes, and liver stellate cells Author Pu Pengyi Zhou 36 Shadan S Yarandi 33 Dongdong Wang 33 Storage: Storage at -20°C. Stable for one year af Storage Buffer:	in components of the intermedia gration, cell motility, maintaini cells, predominantly in astrocyt int in peripheral glia and in non- (PMID: 21219963). bmed ID Journal 260151 J Mol Histol 2137166 PLoS One 2819195 Aging (Alban) ter shipment. 29% glycerol, pH7.3	ate filament network in astrocytes and has beeing mechanical strength, and in mitosis. GFAP is es. GFAP is commonly used as an astrocyte CNS cells, including fibroblasts, chondrocytes, Application IF,WB	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 23935-1-AP (GFAP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-GFAP (IP:23935-1-AP, 4ug; Detection:23935-1-AP 1:20000) with mouse brain tissue lysate 1280 ug.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using GFAP antibody (23935-1-AP) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using GFAP antibody (23935-1-AP) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




WB result of GFAP antibody (23935-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GFAP transfected U-251 cells.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 23935-1-AP (GFAP antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).





Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 23935-1-AP (GFAP antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using GFAP antibody (23935-1-AP) at dilution of 1:200 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).